




Europäisches
Patentamt

European
Patent Office

Office européen
des brevets



Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

03356205.9

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

BEST AVAILABLE COPY



Anmeldung Nr:
Application no.: 03356205.9
Demande no:

Anmeldetag:
Date of filing: 19.12.03
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Bayer CropScience S.A.
16 rue Jean-Marie-Leclair
69009 Lyon
FRANCE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

2-Pyridinylethylbenzamide derivatives

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)

Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

A01N43/400

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PT RO SE SI SK TR LI

Bemerkungen:

Remarks:

Remarques:

See page 1 of the description for the original title

New 2-pyridinylethylbenzamide derivatives


The present invention relates to novel N-[2-(2-pyridinyl)ethyl]benzamide derivatives, their process of preparation, their use as fungicides, particularly in the form of fungicidal compositions, and methods for the control of phytopathogenic fungi of plants using these compounds or their compositions.

International patent application WO 01/11965 discloses a broad family of fungicidal compounds. There is no specific disclosure of N-[2-(2-pyridinyl)ethyl]benzamide derivatives.

It is always of high-interest in the field of agrochemicals to use pesticidal compounds more active than the compounds already known by the man ordinary skilled in the art whereby less compound can be used whilst retaining equivalent efficacy.

We have now found a new family of compounds which show enhanced fungicidal activity over the general known family of such compounds.

Accordingly, the present invention relates to a N-[2-(2-pyridinyl)ethyl]benzamide derivative of general formula (I)



in which :

- n is 1, 2, or 3;
- p is 1, 2, 3 or 4;
- R^a is a C₁-C₆-halogenoalkyl having 1 to 5 halogen atoms;
- each substituent X is chosen, independently of the others, as being a hydrogen atom, a halogen atom, a C₁-C₆-alkyl or a C₁-C₆-halogenoalkyl;
- R¹, R², R³ and R⁴ are chosen, independently of the others as being a hydrogen atom, a halogen atom, a cyano group, a hydroxy group, an amino group, a sulfanyl group, a formyl group, a formyloxy group, a formylamino group, a carboxy group, a carbamoyl group, a N-hydroxycarbamoyl group, a carbamate group, a (hydroxyimino)-C₁-C₆-alkyl group, a C₁-C₆-alkyl, a C₂-C₆-alkenyl, a C₂-C₆-alkynyl,

a C₁-C₆-alkylamino, a di-C₁-C₆-alkylamino, a C₁-C₆-alkoxy, a C₁-C₆-halogenoalkyl having 1 to 5 halogen atoms, a C₁-C₆-halogenoalkoxy having 1 to 5 halogen atoms, a C₁-C₆-alkylsulfanyl, a C₁-C₆-halogenoalkylsulfanyl having 1 to 5 halogen atoms, a C₂-C₆-alkenyloxy, a C₂-C₆-halogenoalkenyloxy having 1 to 5 halogen atoms, a C₃-C₆-alkynyloxy, a C₃-C₆-halogenoalkynyloxy having 1 to 5 halogen atoms, a C₃-C₆-cycloalkyl, a C₃-C₆-halogenocycloalkyl having 1 to 5 halogen atoms, a C₁-C₆-alkylcarbonyl, a C₁-C₆-halogenoalkylcarbonyl having 1 to 5 halogen atoms, a C₁-C₆-alkylcarbamoyl, a di-C₁-C₆-alkylcarbamoyl, a N-C₁-C₆-alkyloxycarbamoyl, a C₁-C₆-alkoxycarbamoyl, a N-C₁-C₆-alkyl-C₁-C₆-alkoxycarbamoyl, a C₁-C₆-alkoxycarbonyl, a C₁-C₆-halogenoalkoxycarbonyl having 1 to 5 halogen atoms, a C₁-C₆-alkylcarbonyloxy, a C₁-C₆-halogenoalkylcarbonyloxy having 1 to 5 halogen atoms, a C₁-C₆-alkylcarbonylamino, a C₁-C₆-halogenoalkylcarbonylamino having 1 to 5 halogen atoms, a C₁-C₆-alkylaminocarbonyloxy, a di-C₁-C₆-alkylaminocarbonyloxy, a C₁-C₆-alkyloxycarbonyloxy, a C₁-C₆-alkylsulphenyl, a C₁-C₆-halogenoalkylsulphenyl having 1 to 5 halogen atoms, a C₁-C₆-alkylsulphinyl, a C₁-C₆-halogenoalkylsulphinyl having 1 to 5 halogen atoms, a C₁-C₆-alkylsulphonyl, a C₁-C₆-halogenoalkylsulphonyl having 1 to 5 halogen atoms, a benzyl, a benzyloxy, a benzylsulfanyl, a benzylsulfinyl, a benzylsulfonyl, a benzylamino, a phenoxy, a phenylsulfanyl, a phenylsulfinyl, a phenylsulfonyl, a phenylamino, a phenylcarbonylamino, a 2,6 dichlorophenyl-carbonylamino group or a phenyl group, with the proviso that when three of the four substituents R¹, R², R³ and R⁴ are a hydrogen atom, then the fourth substituent is not a hydrogen atom;

- R⁵ is chosen as being a hydrogen atom, a cyano group, a formyl group, a hydroxy group, a C₁-C₆-alkyl, a C₁-C₆-halogenoalkyl having 1 to 5 halogen atoms, a C₁-C₆-alkoxy, a C₁-C₆-halogenoalkoxy having 1 to 5 halogen atoms, a C₃-C₆-halogenocycloalkyl having 1 to 5 halogen atoms, a C₂-C₆-alkenyl, a C₂-C₆-alkynyl, a C₁-C₆-alkoxy-C₁-C₆-alkyl, a C₁-C₆-cyanoalkyl, a C₁-C₆-aminoalkyl, a C₁-C₆-alkylamino-C₁-C₆-alkyl, a di-C₁-C₆-alkylamino-C₁-C₆-alkyl, a C₁-C₆-alkylcarbonyl, a C₁-C₆-halogenalkylcarbonyl having 1 to 5 halogen atoms, a C₁-C₆-alkyloxycarbonyl, a C₃-C₇-cycloalkyl, a C₃-C₇-halogenocycloalkyl having 1 to 5 halogen atoms, a C₃-C₇-cycloalkyl-C₁-C₆-alkyl, a C₁-C₆-benzyloxycarbonyl, a C₁-C₆-alkoxy-C₁-C₆-alkylcarbonyl, a C₁-C₆-alkylsulfonyl or a C₁-C₆-halogenoalkylsulfonyl having 1 to 5 halogen atoms;

- Y is the same or different and is a hydrogen atom, a halogen atom, a nitro group, a cyano group, a hydroxy group, an amino group, a sulfanyl group, a pentafluoro-λ⁶-sulfanyl group, a formyl group, a formyloxy group, a formylamino

group, a carboxy group, a C₁-C₈-alkyl, a C₁-C₈-halogenoalkyl having 1 to 5 halogen atoms, a C₂-C₈-alkenyl, a C₂-C₈-alkynyl, a C₁-C₈-alkylamino, a di-C₁-C₈-alkylamino, a C₁-C₈-alkoxy, a C₁-C₈-halogenoalkoxy having 1 to 5 halogen atoms, a C₁-C₈-alkoxy-C₂-C₈-alkenyl, a C₁-C₈-alkylsulfanyl, a C₁-C₈-halogenoalkylsulfanyl having 1 to 5 halogen atoms, a C₁-C₈-alkoxycarbonyl, a C₁-C₈-halogenoalkoxycarbonyl having 1 to 5 halogen atoms, a C₁-C₈-alkylcarbonyloxy, a C₁-C₈-halogenoalkylcarbonyloxy having 1 to 5 halogen atoms, a C₁-C₈-alkylsulphenyl, a C₁-C₈-halogenoalkylsulphenyl having 1 to 5 halogen atoms, a C₁-C₈-alkylsulphinyl, a C₁-C₈-halogenoalkylsulphinyl having 1 to 5 halogen atoms, a C₁-C₈-alkylsulphonyl, a C₁-C₈-halogenoalkylsulphonyl having 1 to 5 halogen atoms or a C₁-C₈-alkylsulfonamide; and

- R^b is a halogen atom, a nitro group, a cyano group, an amino group, a sulfanyl group, a pentafluoro-λ⁶-sulfanyl group, a formyl group, a formyloxy group, a formylamino group, a carboxy group, a C₁-C₆-alkyl, a C₁-C₆-halogenoalkyl having 1 to 5 halogen atoms, a C₂-C₆-alkenyl, a C₂-C₆-alkynyl, a C₁-C₆-alkylamino, a di-C₁-C₆-alkylamino, a C₁-C₆-alkoxy, a C₁-C₆-halogenoalkoxy having 1 to 5 halogen atoms, a C₁-C₆-alkoxy-C₂-C₆-alkenyl, a C₁-C₆-alkylsulfanyl, a C₁-C₆-halogenoalkylsulfanyl having 1 to 5 halogen atoms, a C₁-C₆-alkoxycarbonyl, a C₁-C₆-halogenoalkoxycarbonyl having 1 to 5 halogen atoms, a C₁-C₆-alkylcarbonyloxy, a C₁-C₆-halogenoalkylcarbonyloxy having 1 to 5 halogen atoms, a C₁-C₆-alkylsulphenyl, a C₁-C₆-halogenoalkylsulphenyl having 1 to 5 halogen atoms, a C₁-C₆-alkylsulphinyl, a C₁-C₆-halogenoalkylsulphinyl having 1 to 5 halogen atoms, a C₁-C₆-alkylsulphonyl, a C₁-C₆-halogenoalkylsulphonyl having 1 to 5 halogen atoms or a C₁-C₆-alkylsulfonamide;

as well as its salts, N-oxydes, metallic complexes, metalloidic complexes and optically active isomers.

In the context of the present invention :

- halogen means fluorine, bromine, chlorine or iodine;
- heteroatom means N, O or S;
- carboxy means -C(=O)OH ;
- carbonyl means -C(=O)- ;
- carbamoyl means -C(=O)NH₂ ;
- N-hydroxycarbamoyl means -C(=O)NHOH ;
- an alkyl group, an alkenyl group, and an alkynyl group as well as moieties containing these terms, can be linear or branched.

In the context of the present invention, it has also to be understood that in the case of di-substituted amino and of di-substituted carbamoyl radicals, the two substituents may form together with the nitrogen atom bearing them a saturated heterocyclic ring containing 3 to 7 atoms.

5

According to the present invention, the 2-pyridyl may be substituted in every position by $(X)_n$ and R^a , in which X, R^a and n are as defined above. Preferably, the present invention relates to N-[2-(2-pyridinyl)ethyl]benzamide derivative of general formula (I) in which the different characteristics may be chosen alone or in combination as being :

10

- as regards n, n is 1 or 2. More preferably n is 1.
- as regards X, X is preferably chosen as being a halogen atom. More preferably X is chlorine;
- as regards R^a , R^a is preferably chosen as being $-CF_3$;
- 15 - as regards the positions in which the 2-pyridyl is substituted, the 2-pyridyl is substituted in 3- and/or in 5-position. More preferably, the 2-pyridyl is substituted in 3-position by X and in 5-position by R^a .

Even more preferably, the 2-pyridyl is substituted in 3-position by $-Cl$ and in 5-position by $-CF_3$.

20

According to the present invention, the phenyl is substituted in ortho position by R^b and may be substituted in any other position by $(Y)_p$, in which Y and p are as defined above. Preferably, the present invention relates to N-[2-(2-pyridinyl)ethyl]benzamide derivative of general formula (I) in which the different characteristics may be chosen alone or in combination as being :

25

- as regards R^b , R^b is a halogen atom, a C_1 - C_6 -alkyl, a C_1 - C_6 -alkoxy or a C_1 - C_6 -halogenoalkyl having 1 to 5 halogen atoms;
- as regards p, p is 1;
- as regards Y, Y is a hydrogen atom, a halogen atom or a C_1 - C_6 -alkyl. More preferably, Y is hydrogen.
- 30


According to the present invention, the two carbon atoms and the nitrogen atom of the "ethylamide part" of the compound of formula (I) are respectively substituted by R^1 and R^2 , R^3 and R^4 , and R^5 , at least one the substituents R^1 , R^2 , R^3 and R^4 being different from hydrogen. Preferably, the present invention also relates

35


to N-[2-(2-pyridinyl)ethyl]benzamide derivative of general formula (I) in which the different characteristics may be chosen alone or in combination as being :

- as regards R^1 and R^2 , R^1 and R^2 may be chosen, independently of each other, as being a hydrogen atom, a halogen atom, a cyano group, a hydroxy group, a C_1 - C_6 -alkyl, a C_1 - C_6 -halogenoalkyl having 1 to 5 halogen atoms, a C_2 - C_6 -alkenyl, a C_1 - C_6 -alkoxy, a C_1 - C_6 -alkylsulfanyl, a C_1 - C_6 -alkylsulfenyl, a C_1 - C_6 -alkylsulfinyl, a C_1 - C_6 -alkoxycarbonyl, a C_1 - C_6 -alkylcarbonylamino, a C_1 - C_6 -alkoxycarbonyloxy, a C_1 - C_6 -alkoxycarbonylamino or a phenyl group. More preferably, R^1 and R^2 may be chosen, independently of each other, as being a halogen atom, a C_1 - C_6 -alkyl, a C_1 - C_6 -halogenoalkyl having 1 to 5 halogen atoms or a C_1 - C_6 -alkylcarbonylamino.
- as regards R^3 and R^4 , R^3 and R^4 may be chosen, independently of each other, as being a hydrogen atom, a halogen atom, a cyano group, a C_1 - C_6 -alkyl, a C_1 - C_6 -halogenoalkyl having 1 to 5 halogen atoms, a C_1 - C_6 -alkylcarbonylamino or a phenyl group. More preferably, R^3 and R^4 may be chosen, independently of each other, as being a halogen atom, a C_1 - C_6 -alkyl, a C_1 - C_6 -halogenoalkyl having 1 to 5 halogen atoms or a phenyl group.
- as regards R^5 , R^5 may be chosen as being a hydrogen atom or a C_3 - C_7 -cycloalkyl.

The present invention also relates to a process for the preparation of the compound of general formula (I). Thus, according to a further aspect of the present invention there is provided a process for the preparation of compound of general formula (I) as defined above, which comprises reacting a 2-pyridine derivative of general formula (II) or one of its salt :




in which X, n, R^a , R^1 , R^2 , R^3 , R^4 and R^5 are as defined above;
 30 with a carboxylic acid derivative of the general formula (III)



in which :

- 35 - Y, p and R^b are as defined above; and

- L^2 is a leaving group chosen as being a halogen atom, a hydroxyl group, $-OR^6$, $-OCOR^6$, R^6 being a C_1-C_6 alkyl, a C_1-C_6 haloalkyl, a benzyl, 4-methoxybenzyl, pentafluorophenyl or a group of formula



5


in the presence of a catalyst and, if L^2 is a hydroxyl group, in the presence of a condensing agent.

The process according to the present invention is conducted in the presence of a catalyst. Suitable catalyst may be chosen as being 4-dimethyl-aminopyridine, 1-hydroxy-benzotriazole or dimethylformamide.

In case L^2 is a hydroxy group, the process according to the present invention is conducted in the presence of condensing agent. Suitable condensing agent may be chosen as being acid halide former, such as phosgene, phosphorous tribromide, phosphorous trichloride, phosphorous pentachloride, phosphorous trichloride oxide or thionyl chloride; anhydride former, such as ethyl chloroformate, methyl chloroformate, isopropyl chloroformate, isobutyl chloroformate or methanesulfonyl-chloride; carbodiimides, such as N,N' -dicyclohexylcarbodiimide (DCC) or other customary condensing agents, such as phosphorous pentoxide, polyphosphoric acid, N,N' -carbonyl-diimidazole, 2-ethoxy- N -ethoxycarbonyl-1,2-dihydroquinoline (EEDQ), triphenylphosphine/tetrachloromethane, 4-(4,6-dimethoxy[1.3.5]triazin-2-yl)-4-methylmorpholinium chloride hydrate or bromo-tripyrrolidino-phosphonium-hexafluorophosphate.

25

When R^5 is a hydrogen atom, the above mentioned process for the preparation of compound of general formula (I) may optionally be completed by a further step according to the following reaction scheme :



in which : - $R^1, R^2, R^3, R^4, R^5, R^a, R^b, X, Y, n$ and p are as defined above;

30

- L^5 is a leaving group chosen as being a halogen atom, a 4-methyl phenylsulfonyloxy or a methylsulfonyloxy;


comprising the reaction of a compound of general formula (Id) with a compound of general formula (XXII) to provide a compound of general formula (I).

Depending on the definition of R^1 , R^2 , R^3 , R^4 or R^5 , amine derivatives of general formula (II) may be prepared by different processes. One example (A) of such a process may be when :

- R^a , R^2 , X, n are as defined above;
- R^1 is a hydrogen atom or a C_1 - C_6 alkyl; and
- R^3 , R^4 , R^5 are hydrogen atoms;

then, the amine derivative of general formula (II) may be prepared according to a process which comprises :

- a first step according to reaction scheme A-1 :



in which : - R^a , R^2 , X and n are as defined above;


- R^6 is a C_1 - C_6 alkyl, a C_1 - C_6 haloalkyl, a benzyl, 4-methoxybenzyl or pentafluorophenyl;

- U is a leaving group chosen as being a halogen, a C_1 - C_6 alkylsulfonate or a C_1 - C_6 haloalkylsulfonate;


comprising the arylation of a cyanoacetate derivative of general formula (V) by a pyridine derivative of general formula (IV), to provide a 2-(pyridyl)cyanoacetate derivative of general formula (VI), in the presence of a base, at a temperature of from 0°C to 200°C ;

- a second step according to reaction scheme A-2 :

Scheme A-2




- in which :
- R^a , R^2 , X , n are as defined above;
 - R^6 is a C_1 - C_6 alkyl, a C_1 - C_6 haloalkyl, a benzyl, 4-methoxybenzyl or pentafluorophenyl;
- comprising a basic hydrolysis, an acidic hydrolysis or a displacement by an halide of a compound of general formula (VI) in the same or a different pot to provide, upon heating at a temperature of from 40°C to reflux, a 2-pyridylacetonitrile derivative of general formula (VIIa);
- a third step according to reaction scheme A-3 :



10

- in which :
- R^a , R^2 , X , n are as defined above;
 - R^1 is a C_1 - C_6 alkyl;
 - W is a halogen atom, a C_1 - C_6 alkylsulfonate, a C_1 - C_6 haloalkylsulfonate or a 4-methyl-phenylsulfonate;
- comprising the alkylation of a compound of general formula (VIIa) by a reagent of general formula (XVII) to provide a compound of general formula (VIIb);
- a fourth step according to reaction scheme A-4 :



20


- in which :
- R^a , R^2 , X , n are as defined above;
 - R^1 is a hydrogen atom or a C_1 - C_6 alkyl;
 - L^1 is a leaving group chosen as being a $-OR^6$ group or a $-OCOR^6$ group, R^6 being a C_1 - C_6 alkyl, a C_1 - C_6 haloalkyl, a benzyl, 4-methoxybenzyl or pentafluorophenyl;

- PG represents a protecting group which may be a $-\text{COOR}^6$ group or $-\text{COR}^6$ group, R^6 being a $\text{C}_1\text{-C}_6$ alkyl, a $\text{C}_1\text{-C}_6$ haloalkyl, a benzyl, 4-methoxybenzyl or pentafluorophenyl;

comprising the reduction, by hydrogenation or by an hydride donor, of a compound of general formula (VIIa) or (VIIb), in the presence of a catalyst and in the presence of a compound of general formula (VIII) to produce a compound of general formula (IX), at a temperature of from 0°C to 150°C and under a pressure of from 1 bar and 100 bar;

- a fifth step according to reaction scheme A-5 :

Scheme A-5



in which : - R^a , R^2 , X, n are as defined above;

- R^1 is a hydrogen atom or a $\text{C}_1\text{-C}_6$ alkyl;

- PG represents a protecting group which may be a $-\text{COOR}^6$ group or $-\text{COR}^6$ group, R^6 being a $\text{C}_1\text{-C}_6$ alkyl, a $\text{C}_1\text{-C}_6$ haloalkyl, a benzyl, 4-methoxybenzyl or pentafluorophenyl;

comprising a deprotection reaction, in an acidic or in a basic medium, of a compound of general formula (IX) to provide an amine derivative of general formula (II) or one of its salt.

The first step (step A-1) is conducted in the presence of a base. Preferably, the base will be chosen as being an inorganic or an organic base. Suitable examples of such bases may for example be alkaline earth metal or alkali metal hydrides, hydroxides, amides, alcoholates, carbonates or hydrogen carbonates, acetates or tertiary amines.

The first step (step A-1) according to the present invention is conducted at a temperature of from 0°C to 200°C . Preferably, first step (step A-1) is conducted at a temperature of from 0°C to 120°C , more preferably at a temperature of from 0°C to 80°C .

The first step (step A-1) according to the present invention may be conducted in the presence of a solvent. Preferably, the solvent is chosen as being water, an

organic solvent or a mixture of both. Suitable organic solvents may for example be aliphatic, alicyclic or aromatic solvent.

The first step (step A-1) according to the present invention may also be conducted in the presence of a catalyst. Preferably, the catalyst is chosen as being palladium salts or complexes. More preferably, the catalyst is chosen as being a palladium complex. Suitable palladium complex catalyst may for example be generated directly in the reaction mixture by separately adding to the reaction mixture a palladium salt and a complex ligand. Suitable ligands may for example be bulky phosphines or arsines ligands, such as (R)-(-)-1-[(S)-2-(dicyclohexylphosphino)ferrocenyl]ethyl-dicyclohexylphosphine and its corresponding enantiomer, or a mixture of both; (R)-(-)-1-[(S)-2-(dicyclohexylphosphino)ferrocenyl]ethyl-diphenylphosphine and its corresponding enantiomer, or a mixture of both; (R)-(-)-1-[(S)-2-(diphenylphosphino)ferrocenyl]ethyl-di-t-butylphosphine and its corresponding enantiomer, or a mixture of both; or (R)-(-)-1-[(S)-2-(diphenylphosphino)ferrocenyl]ethyl-dicyclohexylphosphine and its corresponding enantiomer, or a mixture of both.

The fourth step (step A-4) according to the present invention is conducted in the presence of a hydride donor. Preferably, the hydride donor is chosen as being metal or metallloid hydrides such as LiAlH_4 , NaBH_4 , KBH_4 , B_2H_6 .

The fourth step (step A-4) according to the present invention is conducted in the presence of a catalyst. Preferably, the catalyst is chosen as being Co(II)-Chloride, Ni(II)-chloride, ammonia or one of its salt, Palladium on charcoal, Raney Nickel, Raney Cobalt or Platinum.

The fourth step (step A-4) according to the present invention is conducted at a temperature of from 0°C to 150°C . Preferably the temperature is of from 10°C to 120°C . More preferably, the temperature is of from 10°C to 80°C .

The fourth step (step A-4) according to the present invention is conducted under a pressure of from 1 bar to 100 bar. Preferably the pressure is of from 1 bar to 50 bar.

The fourth step (step A-4) according to the present invention may be conducted in the presence of an organic solvent, of water or of a mixture thereof. Preferably, the solvent is chosen as being ether, alcohol, carboxylic acid, or a mixture thereof with water or pure water.


A second example (B) of such a process may be when :

- R^a , R^1 , R^2 , X, n are as defined above; and

- R^3 , R^4 , R^5 are hydrogen atoms;

then, the amine derivative of general formula (II) may be prepared according to a process which comprises :

- a first step according to reaction scheme B-1 :




in which : - R^a , R^1 , R^2 , X and n are as defined above ;

- U is a leaving group chosen as being a halogen atom, a C_1 - C_6 alkylsulfonate or a C_1 - C_6 haloalkylsulfonate;

comprising the arylation of a compound of general formula (Vb) by a pyridine derivative of general formula (IV) to provide a 2-pyridylacetonitrile derivative of general formula (VIIb), in the presence of a base and at a temperature of from -

100°C to 200°C;

- a second step according to reaction scheme B-2 :



in which : - R^a , R^1 , R^2 , X and n are as defined above;

- L^1 is a leaving group chosen as being a $-OR^8$ group or a $-OCOR^8$ group, R^8 being a C_1 - C_6 alkyl, a C_1 - C_6 haloalkyl, a benzyl, 4-methoxybenzyl or pentafluorophenyl;


- PG represents a protecting group which may be a $-COOR^8$ group or $-COR^8$ group, R^8 being a C_1 - C_6 alkyl, a C_1 - C_6 haloalkyl, a benzyl, 4-methoxybenzyl or pentafluorophenyl;

comprising the reduction, by hydrogenation or by an hydride donor, of a compound of general formula (VIIa) or (VIIb), in the presence of a compound of general formula (VIII) to produce a compound of general formula (IX) ;

- a third step according to reaction scheme B-3 :

5

Scheme B-3



in which : - R^a , R^1 , R^2 , X and n are as defined above;

- PG represents a protecting group which may be a $-\text{COOR}^8$ group or $-\text{COR}^8$ group, R^8 being a $\text{C}_1\text{-C}_6$ alkyl, a $\text{C}_1\text{-C}_6$ haloalkyl, a benzyl, 4-methoxybenzyl or pentafluorophenyl;

10

comprising a deprotection reaction, in an acidic or in a basic medium, of a compound of general formula (IX) to provide an amine derivative of general formula (II) or one of its salt.

15

The first step (step B-1) is conducted at a temperature of from -100°C to 200°C . Preferably, first step (step A-1) is conducted at a temperature of from -80°C to 120°C , more preferably at a temperature of from -80°C to 80°C .

The first step (step B-1) according to the present invention is conducted in the presence of a base. Preferably, the base will be chosen as being an inorganic or an organic base. Suitable examples of such bases may for example be alkaline earth metal or alkali metal hydrides, hydroxides, amides, alcoholates, carbonates or hydrogen carbonates, acetates or tertiary amines.

20

The first step (step B-1) according to the present invention may be conducted in the presence of a solvent. Preferably, the solvent is chosen as being water, an organic solvent or a mixture of both. Suitable organic solvents may for example be aliphatic, alicyclic or aromatic solvent.

25

The first step (step B-1) according to the present invention may also be conducted in the presence of a catalyst. Preferably, the catalyst is chosen as being palladium salts or complexes. More preferably, the catalyst is chosen as being a palladium complex. Suitable palladium complex catalyst may for example be generated directly in the reaction mixture by separately adding to the reaction

30

mixture a palladium salt and a complex ligand. Suitable ligands may for example be bulky phosphines or arsines ligands, such as (R)-(-)-1-[(S)-2-(dicyclohexylphosphino)ferrocenyl]ethyldicyclohexylphosphine and its corresponding enantiomer, or a mixture of both; (R)-(-)-1[(S)-2-(dicyclohexylphosphino)ferrocenyl]ethyldiphenylphosphine and its corresponding enantiomer, or a mixture of both; (R)-(-)-1[(S)-2-(diphenylphosphino)ferrocenyl]ethyldi-t-butylphosphine and its corresponding enantiomer, or a mixture of both; or (R)-(-)-1[(S)-2-(diphenylphosphino)ferrocenyl]ethyldicyclohexylphosphine and its corresponding enantiomer, or a mixture of both.


The preferred conditions under which step B-2 is conducted are the same than the preferred conditions under which step A-4 of the above mentioned process A is conducted.

The preferred conditions under which step B-3 is conducted are the same than the preferred conditions under which step A-5 of the above mentioned process A is conducted.

A third example (C) of such a process may be when :

- R^a , R^2 , X, n are as defined above;
 - R^1 is a formylamino, a C_1 - C_8 -alkylcarbonylamino, a C_1 - C_8 -halogenoalkylcarbonylamino having 1 to 5 halogen atoms, a phenylcarbonylamino or a 2,6-dichlorophenylcarbonylamino; and
 - R^3 , R^4 , R^5 are hydrogen atoms;
- then, the amine derivative of general formula (II) may be prepared according to a process comprising :
- a first step according to reaction scheme C-1 :

Scheme C-1




- in which :
- R^a , R^2 , X, n are as defined above;
 - U is a leaving group chosen as being a halogen atom, a C_1 - C_6 alkylsulfonate or a C_1 - C_6 haloalkylsulfonate;

comprising the arylation of a compound of general formula (Vc) by a pyridine derivative of general formula (IV) to provide a 2-pyridylacetonitrile derivative of general formula (VIIc), in the presence of a base and at a temperature of from -100°C to 200°C;

- 5 - a second step according to reaction scheme C-2 :

Scheme C-2




in which R^a , R^2 , X, n are as defined above;

comprising the de-protection, by acidic hydrolysis, of a compound of general formula (VIIc), to produce a compound of general formula (VIId) or one of its salt ;

- 10 - a third step according to reaction scheme C-3 :

Scheme C-3




in which : - R^a , R^2 , X, n are as defined above;

- R^9 is a hydrogen atom, a C_1 - C_6 alkyl group, a C_1 - C_8 -halogenoalkyl having 1 to 5 halogen atoms, a phenyl or 2,6 dichlorophenyl ;

comprising the coupling between a compound of general formula (VIId) and an acyl chloride of general formula (X) to produce a compound of general formula (VIIe).

- 15 - a fourth step according to reaction scheme C-4 :

Scheme C-4



in which : - R^a, R², X, n are as defined above;

- R⁹ is a hydrogen atom, a C₁-C₆ alkyl group, a C₁-C₈-halogenoalkyl having 1 to 5 halogen atoms, a phenyl or 2,6 dichlorophenyl;


- L¹ is a leaving group chosen as being a -OR⁸ group or a OCOR⁸ group, R⁸ being a C₁-C₆ alkyl, a C₁-C₆ haloalkyl, a benzyl, 4-methoxybenzyl or pentafluorophenyl;

- PG represents a protecting group which may be a -COOR⁸ group or -COR⁸ group, R⁸ being a C₁-C₆ alkyl, a C₁-C₆ haloalkyl, a benzyl, 4-methoxybenzyl or pentafluorophenyl;

comprising the reduction, by hydrogenation or by an hydride donor, of a compound of general formula (VIIe), in the presence of a compound of general formula (VIII) to produce a compound of general formula (IXb) ;

- a fifth step according to scheme C-5

Scheme C-5



in which : - R^a, R², X, n are as defined above;

- R⁹ is a hydrogen atom, a C₁-C₆ alkyl group, a C₁-C₈-halogenoalkyl having 1 to 5 halogen atoms, a phenyl or 2,6 dichlorophenyl;

- L¹ is a leaving group chosen as being a -OR⁸ group or a OCOR⁸ group, R⁸ being a C₁-C₆ alkyl, a C₁-C₆ haloalkyl, a benzyl, 4-methoxybenzyl or pentafluorophenyl;


- PG represents a protecting group which may be a $-\text{COOR}^8$ group or $-\text{COR}^8$ group, R^8 being a $\text{C}_1\text{-C}_6$ alkyl, a $\text{C}_1\text{-C}_6$ haloalkyl, a benzyl, 4-methoxybenzyl or pentafluorophenyl;

comprising a deprotection reaction, in an acidic or in a basic medium, of a compound of general formula (IXb) to provide an amine derivative of general formula (II) or one of its salt.

A fourth example (D) of such a process may be when :

- R^a , R^b , R^1 , X, Y, n and p are as defined above;
 - R^2 is a hydrogen atom;
 - R^3 is a $\text{C}_1\text{-C}_6$ alkyl, a $\text{C}_1\text{-C}_6$ haloalkyl, a benzyl or a phenyl; and
 - R^5 is a hydrogen atom, a $\text{C}_1\text{-C}_6$ alkyl, a $\text{C}_1\text{-C}_6$ haloalkyl, a $\text{C}_1\text{-C}_6$ alkoxy or a $\text{C}_3\text{-C}_7$ cycloalkyl;
- then, the amine derivative of general formula (II) may be prepared according to a process comprising :
- a first step according to reaction scheme D-1 :

Scheme D-1



in which : - R^a , R^1 , X and n are as defined above;

- R^3 is a $\text{C}_1\text{-C}_6$ alkyl, a $\text{C}_1\text{-C}_6$ haloalkyl, a benzyl or a phenyl;


- R^6 is a $\text{C}_1\text{-C}_6$ alkyl, a $\text{C}_1\text{-C}_6$ haloalkyl, a benzyl, 4-methoxybenzyl or pentafluorophenyl;

- U is a leaving group chosen as being a halogen, a $\text{C}_1\text{-C}_6$ alkylsulfonate or a $\text{C}_1\text{-C}_6$ haloalkylsulfonate;

comprising the arylation of a ketoacetate derivative of general formula (XI) by a pyridine derivative of general formula (IV), to provide a 2-(pyridyl)ketoacetate derivative of general formula (XII), in the presence of a base, at a temperature of from 0°C to 200°C ;

- a second step according to reaction scheme D-2 :

Scheme D-2



in which : - R^a , R^1 , X, n are as defined above;

- R^2 is a hydrogen atom;


- R^3 is a C_1 - C_6 alkyl, a C_1 - C_6 haloalkyl, a benzyl or a phenyl;

- R^6 is a C_1 - C_6 alkyl, a C_1 - C_6 haloalkyl, a benzyl, 4-methoxybenzyl or pentafluorophenyl;

comprising a basic hydrolysis, an acidic hydrolysis or a displacement by an halide of a compound of general formula (XII) in the same or a different pot to provide, upon heating at a temperature of from 40°C to reflux, a 2-pyridylketone derivative of general formula (XIII);

- a third step according to reaction scheme D-3 :

Scheme D-3



in which : - R^a , R^1 , X and n are as defined above;

- R^2 is a hydrogen atom;

- R^3 is a C_1 - C_6 alkyl, a C_1 - C_6 haloalkyl, a benzyl or a phenyl;


- R^5 is a hydrogen atom, a C_1 - C_6 alkyl, a C_1 - C_6 haloalkyl, a

C_1 - C_6 alkoxy or a C_3 - C_7 cycloalkyl;

comprising the reaction of a compound of general formula (XIII) with an amine of formula R^5 -NH₂ to provide an imine derivative of general formula (XIV);

- a fourth step according to scheme D-4 :

Scheme D-4



in which : - R^a , R^1 , X and n are as defined above;

- R^2 is a hydrogen atom;

- R^3 is a C_1 - C_6 alkyl, a C_1 - C_6 haloalkyl, a benzyl, a phenyl;

- R^5 is a hydrogen atom, a C_1 - C_6 alkyl, a C_1 - C_6 haloalkyl, a

C_1 - C_6 alkoxy or a C_3 - C_7 cycloalkyl;

comprising the reduction of an imine derivative of general formula (XIV) by
hydrogenation or by an hydride donor, in the same or a different pot to provide an
amine derivative of general formula (II) or one of its salt.

A fifth example (E) of such a process may be when :

- R^a , R^b , R^1 , R^2 , X, Y, n and p are as defined above;

- R^3 is a hydrogen atom

- R^4 is a C_1 - C_6 alkyl, a C_1 - C_6 haloalkyl, a benzyl, a phenyl; and


- R^5 is a hydrogen atom, a C_1 - C_6 alkyl, a C_1 - C_6 haloalkyl, a C_1 - C_6 alkoxy or a C_3 - C_7 cycloalkyl;

then, the amine derivative of general formula (II) may be prepared according to a

process comprising :

- a first step according to reaction scheme E-1 :

Scheme E-1



in which : - R^a , R^1 , R^2 , X and n are as defined above;

- R^4 is a C_1 - C_6 alkyl, a C_1 - C_6 haloalkyl, a benzyl or a phenyl;


- R^6 is a C_1 - C_6 alkyl, a C_1 - C_6 haloalkyl, a benzyl, 4-methoxybenzyl or pentafluorophenyl;

- U is a leaving group chosen as being a halogen, a C₁-C₆ alkylsulfonate or a C₁-C₆ haloalkylsulfonate;

comprising the arylation of a ketone derivative of general formula (XIV) by a pyridine derivative of general formula (IV), to provide a 2-(pyridyl)ketone derivative of general formula (XIIIb), in the presence of a base, at a temperature of from 0°C to 200°C;

- a second step according to reaction scheme E-2 :

Scheme E-2




in which :
 - R^a, R¹, R², X and n are as defined above;
 - R⁴ is a C₁-C₆ alkyl, a C₁-C₆ haloalkyl, a benzyl or a phenyl;
 - R⁵ is a hydrogen atom, a C₁-C₆ alkyl, a C₁-C₆ haloalkyl, a C₁-C₆ alkoxy or a C₃-C₇ cycloalkyl;

comprising the reaction of a compound of general formula (XIIIb) with an amine of formula R⁵-NH₂ to provide an imine derivative of general formula (XIVb);

- a third step according to scheme E-3 :

Scheme E-3




in which :
 - R^a, R¹, R², X and n are as defined above;
 - R³ is a C₁-C₆ alkyl, a C₁-C₆ haloalkyl, a benzyl or a phenyl;
 - R⁴ is a hydrogen atom;
 - R⁵ is a hydrogen atom, a C₁-C₆ alkyl, a C₁-C₆ haloalkyl, a C₁-C₆ alkoxy or a C₃-C₇ cycloalkyl;

comprising the reduction of an imine derivative of general formula (XIVb) by hydrogenation or by an hydride donor, in the same or a different pot to provide an amine derivative of general formula (II) or one of its salt.

5 A sixth example (F) of such a process may be when :

- R^a , R^2 , X, n are as defined above;
 - R^1 is a cyano group, a hydroxy group, a C_2 - C_6 -alkynyl, a C_1 - C_6 -alkylamino, a di- C_1 - C_6 -alkylamino, a C_1 - C_6 -alkoxy, a C_1 - C_6 -halogenoalkoxy having 1 to 5 halogen atoms, a C_1 - C_6 -alkylsulfanyl, a C_1 - C_6 -halogenoalkylsulfanyl having 1 to 5 halogen atoms, a C_2 - C_6 -alkenyloxy, a C_2 - C_6 -halogenoalkenyloxy having 1 to 5 halogen atoms, a C_3 - C_6 -alkynyloxy, a C_3 - C_6 -halogenoalkynyloxy having 1 to 5 halogen atoms, a benzyloxy, a benzylsulfanyl, a benzylamino, a phenoxy, a phenylsulfanyl, a phenylamino, or a phenyl sulphanyl group, a C_1 - C_6 -alkylcarbonyloxy, a C_1 - C_6 -halogenoalkylcarbonyloxy having 1 to 5 halogen atoms; and
 - 15 - R^3 and R^4 are hydrogen atoms;
- then, the amine derivative of general formula (II) may be prepared according to a process comprising :

- a first step according to reaction scheme F-1 :



20

in which : - R^a , R^2 , X and n are as defined above;

- R^6 is a C_1 - C_6 alkyl, a C_1 - C_6 haloalkyl, a benzyl, 4-methoxybenzyl or pentafluorophenyl;


25 - U is a leaving group chosen as being a halogen, a C_1 - C_6 alkylsulfonate or a C_1 - C_6 haloalkylsulfonate;

comprising the arylation of a cyanoacetate derivative of general formula (V) by a pyridine derivative of general formula (IV), to provide a 2-(pyridyl)cyanoacetate derivative of general formula (VI), in the presence of a base, at a temperature of from

30 0°C to 200°C ;

- a second step according to reaction scheme F-2 :

Scheme F-2



in which : - R^a , R^2 , X, n are as defined above;


- R^1 is a hydrogen atom;

- R^6 is a C_1 - C_6 alkyl, a C_1 - C_6 haloalkyl, a benzyl, 4-methoxybenzyl or pentafluorophenyl;

comprising a basic hydrolysis, an acidic hydrolysis or a displacement by an halide of a compound of general formula (VI) in the same or a different pot to provide, upon heating at a temperature of from 40°C to reflux, a 2-pyridylacetonitrile derivative of general formula (VIIa);

- a third step according to reaction scheme F-3 :

Scheme F-3




in which : - R^a , R^2 , X, n are as defined above;

- W is a halogen atom;

comprising the halogenation of a compound of general formula (VIIa) to provide a compound of general formula (VIIf);


- a fourth step according to reaction scheme F-4 :

Scheme F-4



- in which :
- R^a , R^2 , X , n are as defined above;
 - W is a halogen atom;
 - R^1 is a cyano group, a hydroxy group, a C_2 - C_6 -alkynyl, a C_1 - C_6 -alkylamino, a di- C_1 - C_6 -alkylamino, a C_1 - C_6 -alkoxy, a C_1 - C_6 -halogenoalkoxy having 1 to 5 halogen atoms, a C_1 - C_6 -alkylsulfanyl, a C_1 - C_6 -halogenoalkylsulfanyl having 1 to 5 halogen atoms, a C_2 - C_6 -alkenyloxy, a C_2 - C_6 -halogenoalkenyloxy having 1 to 5 halogen atoms, a C_3 - C_6 -alkynyloxy, a C_3 - C_6 -halogenoalkynyloxy having 1 to 5 halogen atoms, a benzyloxy, a benzylsulfanyl, a benzylamino, a phenoxy, a phenylsulfanyl, a phenylamino, or a phenyl sulphanyl group, a C_1 - C_6 -alkylcarbonyloxy or a C_1 - C_6 -halogenoalkylcarbonyloxy having 1 to 5 halogen atoms; comprising the nucleophilic substitution, in the presence of a base, of a compound of general formula (VIIf) in the presence of a compound of general formula (XV) to produce a compound of general formula (VIIb), at a temperature of from -78°C to 150°C ,
 - a fifth step according to reaction scheme F-5 :

Scheme F-5




- in which :
- R^a , R^2 , X , n are as defined above;
 - R^1 is a cyano group, a hydroxy group, a C_2 - C_6 -alkynyl, a C_1 - C_6 -alkylamino, a di- C_1 - C_6 -alkylamino, a C_1 - C_6 -alkoxy, a C_1 - C_6 -halogenoalkoxy having 1 to 5 halogen atoms, a C_1 - C_6 -alkylsulfanyl, a C_1 - C_6 -halogenoalkylsulfanyl having 1 to 5 halogen atoms, a C_2 - C_6 -alkenyloxy, a C_2 - C_6 -halogenoalkenyloxy having 1 to 5 halogen atoms, a C_3 - C_6 -alkynyloxy, a C_3 - C_6 -halogenoalkynyloxy having 1 to 5 halogen atoms, a benzyloxy, a benzylsulfanyl, a benzylamino, a phenoxy, a phenylsulfanyl, a phenylamino, or a phenyl sulphanyl group, a C_1 - C_6 -alkylcarbonyloxy or a C_1 - C_6 -halogenoalkylcarbonyloxy having 1 to 5 halogen atoms;
 - L^1 is a leaving group chosen as being a $-OR^8$ group or a $-OCOR^8$ group, R^8 being a C_1 - C_6 alkyl, a C_1 - C_6 haloalkyl, a benzyl, 4-methoxybenzyl or pentafluorophenyl;

- PG represents a protecting group which may be a $-\text{COOR}^8$ group or $-\text{COR}^8$ group, R^8 being a $\text{C}_1\text{-C}_6$ alkyl, a $\text{C}_1\text{-C}_6$ haloalkyl, a benzyl, 4-methoxybenzyl or pentafluorophenyl;

comprising the reduction, by hydrogenation or by an hydride donor, of a compound of general formula (VIIb), in the presence of a catalyst and in the presence of a compound of general formula (VIII) to produce a compound of general formula (IX), at a temperature of from 0°C to 150°C and under a pressure of from 1 bar and 100 bar;

- a sixth step according to reaction scheme F-6 :

Scheme F-6



in which : - R^a , R^2 , X , n are as defined above;

- R^1 is a cyano group, a hydroxy group, a $\text{C}_2\text{-C}_6$ -alkynyl, a $\text{C}_1\text{-C}_6$ -alkylamino, a di- $\text{C}_1\text{-C}_6$ -alkylamino, a $\text{C}_1\text{-C}_6$ -alkoxy, a $\text{C}_1\text{-C}_6$ -halogenoalkoxy having 1 to 5 halogen atoms, a $\text{C}_1\text{-C}_6$ -alkylsulfanyl, a $\text{C}_1\text{-C}_6$ -halogenoalkylsulfanyl having 1 to 5 halogen atoms, a $\text{C}_2\text{-C}_6$ -alkenyloxy, a $\text{C}_2\text{-C}_6$ -halogenoalkenyloxy having 1 to 5 halogen atoms, a $\text{C}_3\text{-C}_6$ -alkynyloxy, a $\text{C}_3\text{-C}_6$ -halogenoalkynyloxy having 1 to 5 halogen atoms, a benzyloxy, a benzylsulfanyl, a benzylamino, a phenoxy, a phenylsulfanyl, a phenylamino, or a phenyl sulphanyl group, a $\text{C}_1\text{-C}_6$ -alkylcarbonyloxy or a $\text{C}_1\text{-C}_6$ -halogenoalkylcarbonyloxy having 1 to 5 halogen atoms;

- PG represents a protecting group which may be a $-\text{COOR}^8$ group or $-\text{COR}^8$ group, R^8 being a $\text{C}_1\text{-C}_6$ alkyl, a $\text{C}_1\text{-C}_6$ haloalkyl, a benzyl, 4-methoxybenzyl or pentafluorophenyl; comprising a deprotection reaction, in an acidic or in a basic medium, of a compound of general formula (IX) to provide an amine derivative of general formula (II) or one of its salt.


A seventh example (G) of such a process may be when :

- R^a , X , n are as defined above;
- R^1 is a hydroxy group; and
- R^3 , R^2 , and R^4 are hydrogen atoms;

then, the amine derivative of general formula (II) may be prepared according to a process comprising :

- a first step according to reaction scheme G-1 :

Scheme G-1




in which :
 - R^a , X and n are as defined above;
 - Z is a halogen atom;

comprising the addition of a methyl magnesium halogenide of general formula (XVII) on a pyridine derivative of general formula (XVI) to provide a compound of general formula (XVIII);

- a second step according to reaction scheme G-2 :


Scheme G-2



in which :
 - R^a , X and n are as defined above;
 - Z is a halogen atom;


comprising the halogenation of a compound of general formula (XVIII) into a compound of general formula (XIX) by use of a halogenating agent such as chlorine, bromine, hypochlorite ions, hypobromite ions, trichloride ions, tribromide ions, N-chloro imides, N-chloro amides, N-chloro amines, N-bromo imides, N-bromo amides or N-bromo amines;

- a third step according to reaction scheme G-3 :


Scheme G-3

in which : - R^a , X and n are as defined above;
 - Z is a halogen atom;

- comprising the nucleophilic substitution of a compound of general formula (XIX) by
 5 a phthalimide salt to produce a compound of general formula (XX);
 - a fourth step according to reaction scheme G-4 :

Scheme G-4

- in which R^a , X and n are as defined above;
 10 comprising the reduction of a compound of general formula (XX) by a hydride
 donor to produce a compound of general formula (XXI);
 - a fifth step according to reaction scheme G-5 :

Scheme G-5

- 15 in which R^a , X and n are as defined above;

comprising the de-protection of a compound of general formula (XXI) by reacting it with hydrazine hydrate or a hydrazine salt to provide an amine derivative of general formula (II) or one of its salt;

5 Compounds according to the present invention can be prepared according to the above described processes. It will nevertheless be understood that, on the basis of his general knowledge and of available publications, the skilled worker will be able to adapt these processes according to the specifics of each of the compounds which it is desired to synthesise.

10 The present invention also relates to a fungicidal composition comprising an effective amount of an active material of general formula (I). Thus, according to the present invention, there is provided a fungicidal composition comprising, as an active ingredient, an effective amount of a compound of general formula (I) as defined
15 above and an agriculturally acceptable support, carrier or filler.

 In the present specification, the term "support" denotes a natural or synthetic, organic or inorganic material with which the active material is combined to make it easier to apply, notably to the parts of the plant. This support is thus generally inert and should be agriculturally acceptable. The support may be a solid or a liquid.
20 Examples of suitable supports include clays, natural or synthetic silicates, silica, resins, waxes, solid fertilisers, water, alcohols, in particular butanol, organic solvents, mineral and plant oils and derivatives thereof. Mixtures of such supports may also be used.

 The composition may also comprise additional components. In particular, the
25 composition may further comprise a surfactant. The surfactant can be an emulsifier, a dispersing agent or a wetting agent of ionic or non-ionic type or a mixture of such surfactants. Mention may be made, for example, of polyacrylic acid salts, lignosulphonic acid salts, phenolsulphonic or naphthalenesulphonic acid salts, polycondensates of ethylene oxide with fatty alcohols or with fatty acids or with fatty
30 amines, substituted phenols (in particular alkylphenols or arylphenols), salts of sulphosuccinic acid esters, taurine derivatives (in particular alkyl taurates), phosphoric esters of polyoxyethylated alcohols or phenols, fatty acid esters of polyols, and derivatives of the above compounds containing sulphate, sulphonate and phosphate functions. The presence of at least one surfactant is generally essential
35 when the active material and/or the inert support are water-insoluble and when the

vector agent for the application is water. Preferably, surfactant content may be comprised between 5% and 40% by weight of the composition.

Optionally, additional components may also be included, e.g. protective colloids, adhesives, thickeners, thixotropic agents, penetration agents, stabilisers, sequestering agents. More generally, the active materials can be combined with any solid or liquid additive, which complies with the usual formulation techniques.

In general, the composition according to the invention may contain from 0.05 to 99% (by weight) of active material, preferably 10 to 70% by weight.

Compositions according to the present invention can be used in various forms such as aerosol dispenser, capsule suspension, cold fogging concentrate, dustable powder, emulsifiable concentrate, emulsion oil in water, emulsion water in oil, encapsulated granule, fine granule, flowable concentrate for seed treatment, gas (under pressure), gas generating product, granule, hot fogging concentrate, macrogranule, microgranule, oil dispersible powder, oil miscible flowable concentrate, oil miscible liquid, paste, plant rodlet, powder for dry seed treatment, seed coated with a pesticide, soluble concentrate, soluble powder, solution for seed treatment, suspension concentrate (flowable concentrate), ultra low volume (ulv) liquid, ultra low volume (ulv) suspension, water dispersible granules or tablets, water dispersible powder for slurry treatment, water soluble granules or tablets, water soluble powder for seed treatment and wettable powder.

These compositions include not only compositions which are ready to be applied to the plant or seed to be treated by means of a suitable device, such as a spraying or dusting device, but also concentrated commercial compositions which must be diluted before application to the crop.

The compounds of the invention can also be mixed with one or more insecticides, fungicides, bactericides, attractant acaricides or pheromones or other compounds with biological activity. The mixtures thus obtained have a broadened spectrum of activity. The mixtures with other fungicides are particularly advantageous.

The fungicidal compositions of the present invention can be used to curatively or preventively control the phytopathogenic fungi of crops. Thus, according to a further aspect of the present invention, there is provided a method for curatively or preventively controlling the phytopathogenic fungi of crops characterised in that a fungicidal composition as hereinbefore defined is applied to the seed, the plant and/or to

the fruit of the plant or to the soil in which the plant is growing or in which it is desired to grow.

The composition as used against phytopathogenic fungi of crops comprises an effective and non-phytotoxic amount of an active material of general formula (I).

5 The expression "effective and non-phytotoxic amount" means an amount of composition according to the invention which is sufficient to control or destroy the fungi present or liable to appear on the crops, and which does not entail any appreciable symptom of phytotoxicity for the said crops. Such an amount can vary within a wide range depending on the fungus to be controlled, the type of crop, the climatic conditions
10 and the compounds included in the fungicidal composition according to the invention.

This amount can be determined by systematic field trials, which are within the capabilities of a person skilled in the art.

The method of treatment according to the present invention is useful to treat propagation material such as tubers or rhizomes, but also seeds, seedlings or
15 seedlings pricking out and plants or plants pricking out. This method of treatment can also be useful to treat roots. The method of treatment according to the present invention can also be useful to treat the overground parts of the plant such as trunks, stems or stalks, leaves, flowers and fruits of the concerned plant.

Among the plants that can be protected by the method according to the
20 invention, mention may be made of cotton; flax; vine; fruit crops such as *Rosaceae* sp. (for instance pip fruits such as apples and pears, but also stone fruits such as apricots, almonds and peaches), *Ribesioideae* sp., *Juglandaceae* sp., *Betulaceae* sp., *Anacardiaceae* sp., *Fagaceae* sp., *Moraceae* sp., *Oleaceae* sp., *Actinidaceae* sp., *Lauraceae* sp., *Musaceae* sp. (for instance banana trees and plantains), *Rubiaceae* sp.,
25 *Theaceae* sp., *Sterculiaceae* sp., *Rutaceae* sp. (for instance lemons, oranges and grapefruits); leguminous crops such as *Solanaceae* sp. (for instance tomatoes), *Liliaceae* sp., *Asteraceae* sp. (for instance lettuces), *Umbelliferae* sp., *Cruciferae* sp., *Chenopodiaceae* sp., *Cucurbitaceae* sp., *Papilionaceae* sp. (for instance peas), *Rosaceae* sp. (for instance strawberries); big crops such as *Graminae* sp. (for
30 instance maize, cereals such as wheat, rice, barley and triticale), *Asteraceae* sp. (for instance sunflower), *Cruciferae* sp. (for instance colza), *Papilionaceae* sp. (for instance soja), *Solanaceae* sp. (for instance potatoes), *Chenopodiaceae* sp. (for instance beetroots); horticultural and forest crops; as well as genetically modified homologues of these crops.

35 Among the plants and the possible diseases of these plants protected by the method according to the present invention, mention may be made of :

- wheat, as regards controlling the following seed diseases: fusaria (*Microdochium nivale* and *Fusarium roseum*), stinking smut (*Tilletia caries*, *Tilletia controversa* or *Tilletia indica*), septoria disease (*Septoria nodorum*) and loose smut;
- wheat, as regards controlling the following diseases of the aerial parts of the
 5 plant: cereal eyespot (*Tapesia yallundae*, *Tapesia acuiformis*), take-all (*Gaeumannomyces graminis*), foot blight (*F. culmorum*, *F. graminearum*), black speck (*Rhizoctonia cerealis*), powdery mildew (*Erysiphe graminis forma specie tritici*), rusts (*Puccinia striiformis* and *Puccinia recondita*) and septoria diseases (*Septoria tritici* and *Septoria nodorum*);
- 10 - wheat and barley, as regards controlling bacterial and viral diseases, for example barley yellow mosaic;
- barley, as regards controlling the following seed diseases: net blotch (*Pyrenophora graminea*, *Pyrenophora teres* and *Cochliobolus sativus*), loose smut (*Ustilago nuda*) and fusaria (*Microdochium nivale* and *Fusarium roseum*);
- 15 - barley, as regards controlling the following diseases of the aerial parts of the plant: cereal eyespot (*Tapesia yallundae*), net blotch (*Pyrenophora teres* and *Cochliobolus sativus*), powdery mildew (*Erysiphe graminis forma specie hordei*), dwarf leaf rust (*Puccinia hordei*) and leaf blotch (*Rhynchosporium secalis*);
- potato, as regards controlling tuber diseases (in particular *Helminthosporium solani*, *Phoma tuberosa*, *Rhizoctonia solani*, *Fusarium solani*), mildew (*Phytophthora infestans*) and certain viruses (virus Y);
- 20 - potato, as regards controlling the following foliage diseases: early blight (*Alternaria solani*), mildew (*Phytophthora infestans*);
- cotton, as regards controlling the following diseases of young plants grown
 25 from seeds: damping-off and collar rot (*Rhizoctonia solani*, *Fusarium oxysporum*) and black root rot (*Thielaviopsis basicola*);
- protein yielding crops, for example peas, as regards controlling the following seed diseases: anthracnose (*Ascochyta pisi*, *Mycosphaerella pinodes*), fusaria (*Fusarium oxysporum*), grey mould (*Botrytis cinerea*) and mildew
 30 (*Peronospora pisi*);
- oil-bearing crops, for example rape, as regards controlling the following seed diseases: *Phoma lingam*, *Alternaria brassicae* and *Sclerotinia sclerotiorum*;
- corn, as regards controlling seed diseases: (*Rhizopus* sp., *Penicillium* sp., *Trichoderma* sp., *Aspergillus* sp., and *Gibberella fujikuroi*);

- flax, as regards controlling the seed disease: *Alternaria linicola*;
- forest trees, as regards controlling damping-off (*Fusarium oxysporum*, *Rhizoctonia solani*);
- rice, as regards controlling the following diseases of the aerial parts: blast disease (*Magnaporthe grisea*), bordered sheath spot (*Rhizoctonia solani*);
- leguminous crops, as regards controlling the following diseases of seeds or of young plants grown from seeds: damping-off and collar rot (*Fusarium oxysporum*, *Fusarium roseum*, *Rhizoctonia solani*, *Pythium sp.*);
- leguminous crops, as regards controlling the following diseases of the aerial parts: grey mould (*Botrytis sp.*), powdery mildews (in particular *Erysiphe cichoracearum*, *Sphaerotheca fuliginea* and *Leveillula taurica*), fusaria (*Fusarium oxysporum*, *Fusarium roseum*), leaf spot (*Cladosporium sp.*), alternaria leaf spot (*Alternaria sp.*), anthracnose (*Colletotrichum sp.*), septoria leaf spot (*Septoria sp.*), black speck (*Rhizoctonia solani*), mildews (for example *Bremia lactucae*, *Peronospora sp.*, *Pseudoperonospora sp.*, *Phytophthora sp.*);
- fruit trees, as regards diseases of the aerial parts: monilia disease (*Monilia fructigenae*, *M. laxa*), scab (*Venturia inaequalis*), powdery mildew (*Podosphaera leucotricha*);
- vine, as regards diseases of the foliage: in particular grey mould (*Botrytis cinerea*), powdery mildew (*Uncinula necator*), black rot (*Guignardia biwelli*) and mildew (*Plasmopara viticola*);
- beetroot, as regards the following diseases of the aerial parts: cercospora blight (*Cercospora beticola*), powdery mildew (*Erysiphe beticola*), leaf spot (*Ramularia beticola*).

25

The fungicide composition according to the present invention may also be used against fungal diseases liable to grow on or inside timber. The term "timber" means all types of species of wood, and all types of working of this wood intended for construction, for example solid wood, high-density wood, laminated wood, and plywood. The method for treating timber according to the invention mainly consists in contacting one or more compounds of the present invention, or a composition according to the invention; this includes for example direct application, spraying, dipping, injection or any other suitable means.

35


The dose of active material usually applied in the treatment according to the present invention is generally and advantageously between 10 and 800 g/ha, preferably

between 50 and 300 g/ha for applications in foliar treatment. The dose of active substance applied is generally and advantageously between 2 and 200 g per 100 kg of seed, preferably between 3 and 150 g per 100 kg of seed in the case of seed treatment. It is clearly understood that the doses indicated above are given as illustrative examples of the invention. A person skilled in the art will know how to adapt the application doses according to the nature of the crop to be treated.

The fungicidal composition according to the present invention may also be used in the treatment of genetically modified organisms with the compounds according to the invention or the agrochemical compositions according to the invention. Genetically modified plants are plants into whose genome a heterologous gene encoding a protein of interest has been stably integrated. The expression "heterologous gene encoding a protein of interest" essentially means genes which give the transformed plant new agronomic properties, or genes for improving the agronomic quality of the transformed plant.

The compositions according to the present invention may also be used for the preparation of composition useful to curatively or preventively treat human and animal fungal diseases such as, for example, mycoses, dermatoses, trichophyton diseases and candidiases or diseases caused by *Aspergillus spp.*, for example *Aspergillus fumigatus*.

The aspects of the present invention will now be illustrated with reference to the following tables of compounds and examples. The following Tables A to V illustrate in a non-limiting manner examples of fungicidal compounds according to the present invention. In the following Examples, M+1 (or M-1) means the molecular ion peak, plus or minus 1 a.m.u. (atomic mass units) respectively, as observed in mass spectroscopy.



Compound n°	R ^a	X ¹	X ²	X ³	R ¹	R ²	R ³	R ⁴	R ⁵	R ^b	Y ¹	Y ²	Y ³	Y ⁴	M+1
1	CF ₃	Cl	H	H	Me	H	H	H	H	Cl	H	H	H	Cl	411
2	CF ₃	Cl	H	H	Me	H	H	H	H	Cl	H	H	Cl	H	411
3	CF ₃	Cl	H	H	Me	H	H	H	H	Cl	H	Cl	H	H	411
4	CF ₃	Cl	H	H	Me	H	H	H	H	OMe	H	H	H	H	373
5	CF ₃	Cl	H	H	Me	H	H	H	H	OMe	H	H	H	OMe	403
6	CF ₃	Cl	H	H	Me	H	H	H	H	OMe	OMe	H	H	H	403
7	CF ₃	Cl	H	H	Me	H	H	H	H	Me	H	H	H	H	357
8	CF ₃	Cl	H	H	Me	H	H	H	H	Me	H	H	H	Me	371

9	CF ₃	Cl	H	H	H	Me	H	H	H	H	Me	H	H	Me	H	371
10	CF ₃	Cl	H	H	H	Me	H	H	H	H	Me	Me	H	H	H	371
11	CF ₃	Cl	H	H	H	Me	H	H	H	H	F	H	H	H	Cl	395
12	CF ₃	Cl	H	H	H	Me	H	H	H	H	Me	H	H	H	Cl	391
13	CF ₃	Cl	H	H	H	Me	H	H	H	H	Me	H	H	H	F	375
14	CF ₃	Cl	H	H	H	Me	H	H	H	H	F	H	H	Cl	H	395
15	CF ₃	Cl	H	H	H	Me	H	H	H	H	Me	⁴ H	H	H	OH	373
16	CF ₃	Cl	H	H	H	Et	H	H	H	H	Cl	H	H	H	Cl	427
17	CF ₃	Cl	H	H	H	Et	H	H	H	H	NMe	H	H	H	H	425
18	CF ₃	Cl	H	H	H	Et	H	H	H	H	Cl	H	Cl	H	H	427
19	CF ₃	Cl	H	H	H	Et	H	H	H	H	NMe	H	H	H	H	386
20	CF ₃	Cl	H	H	H	Et	H	H	H	H	OMe	H	H	H	H	387
21	CF ₃	Cl	H	H	H	Et	H	H	H	H	OMe	H	H	H	OMe	417
22	CF ₃	Cl	H	H	H	Et	H	H	H	H	OMe	OMe	H	H	H	417

23	CF ₃	Cl	H	H	H	H	H	H	H	Me	H	H	H	H	H	371
24	CF ₃	Cl	H	H	H	H	H	H	H	Me	H	H	H	H	Me	386
25	CF ₃	Cl	H	H	H	H	H	H	H	Me	H	H	H	Me	H	385
26	CF ₃	Cl	H	H	H	H	H	H	H	F	H	H	H	H	Cl	409
27	CF ₃	Cl	H	H	H	H	H	H	H	Me	H	H	H	H	Cl	405
28	CF ₃	Cl	H	H	H	H	H	H	H	Me	H	H	H	H	F	389
29	CF ₃	Cl	H	H	H	H	H	H	H	F	H	Cl	H	H	H	409
30	CF ₃	Cl	H	H	H	H	H	H	H	Me	H	H	H	H	OH	387
31	CF ₃	Cl	H	H	H	H	H	H	H	Me	H	Me	H	H	H	385
32	CF ₃	Cl	H	H	H	H	H	H	H	CF ₃	H	H	H	H	H	411
33	CF ₃	Cl	H	H	H	H	H	Me	H	CF ₃	H	H	H	H	H	425
34	CF ₃	Cl	H	H	H	H	Me	Et	H	CF ₃	H	H	H	H	H	439
35	CF ₃	Cl	H	H	H	Pr	Pr	Pr	H	CF ₃	H	H	H	H	H	481
36	CF ₃	Cl	H	H	H	Pr	Pr	Pr	H	I	H	H	H	H	H	539

37	CF ₃	Cl	H	H	H	Pr	H	H	H	H	Br	H	H	H	H	490
38	CF ₃	Cl	H	H	H	H	Me	H	H	H	CF ₃	H	H	H	H	411
39	CF ₃	Cl	H	H	H	Et	H	H	H	H	CF ₃	H	H	H	H	425
40	CF ₃	Cl	H	H	H	Et	H	H	H	H	Br	H	H	H	H	437
41	CF ₃	Cl	H	H	H	Et	H	H	H	H	CHF ₂	H	H	H	H	407
42	CF ₃	Cl	H	H	H	Pr	H	H	H	H	CF ₃	H	H	H	H	439
43	CF ₃	Cl	H	H	H	Pr	H	H	H	H	Br	H	H	H	H	451
44	CF ₃	Cl	H	H	H	Pr	H	H	H	H	CHF ₂	H	H	H	H	421
45	CF ₃	Cl	H	H	H	OH	H	H	H	H	CF ₃	H	H	H	H	413
46	CF ₃	Cl	H	H	H	Me	H	H	H	H	CHF ₂	H	H	H	H	407
47	CF ₃	Cl	H	H	H	H	Et	H	H	H	CF ₃	H	H	H	H	425
48	CF ₃	Cl	H	H	H	NHCOMe	H	H	H	H	CF ₃	H	H	H	H	454
49	CF ₃	Cl	H	H	H	H	H	H	Ph	H	CF ₃	H	H	H	H	473

Examples of process for the preparation of the compound of general formula (I)

5 Example 1 : Preparation of N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]butyl}-2-(trifluoromethyl)benzamide (compound 39)

100 mg of 2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-1-butanamine (0.0004 mol), 58 μ L of 2-(trifluoromethyl)benzoic acid (0.0004 mol), 0.109 g of 4-(4,6-dimethoxy[1.3.5]triazin-2-yl)-4-methylmorpholinium chloride hydrate (0.0004 mol) are stirred in 2 mL of ethanol at room temperature overnight.

The reaction mixture is concentrated to dryness and purified on silica to yield to 64 mg of N-(1-{[3-chloro-5-(trifluoromethyl)-2-pyridinyl]methyl}propyl)-2-(trifluoromethyl)benzamide (57 %).

Mass spectrum: $[M+1] = 425$.

15

Example 2 : Preparation of N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-2-hydroxyethyl}-2-(trifluoromethyl)benzamide (compound 45)

To 50 mg (0.21 mmol) of 2-amino-1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethanol in solution in 3 ml of acetonitrile, was successfully added 0.044 ml (0.21 mmol) of 2-trifluoromethylbenzoyl chloride and 37 mg (0.21 mmol) of potassium carbonate. The reaction mixture was stirred at room temperature for 18 hours. The reaction mixture was poured into aqueous potassium carbonate and the aqueous phase was extracted with ethyl acetate (2 x 50 ml).

25 The organic phase was washed with brine and dried over magnesium sulphate. The solvent was evaporated under reduced pressure to give pure N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-2-hydroxyethyl}-2-(trifluoromethyl)benzamide: 43 mg (43%).

Mass spectrum: $[M+1] = 413$.

30

Example 3 : Preparation of N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-2-methylpropyl}-2-(difluoromethyl)benzamide (compound 46)

104 mg of ammonium (0.00036 mol), 40 μ L of triethylamine, 62 mg of 2-difluoromethylbenzoic acid, 0.13 g of 4-(4,6-dimethoxy[1.3.5]triazin-2-yl)-4-

35

methylmorpholinium chloride hydrate (0.00054 mol) are stirred in 2 mL of methanol at room temperature overnight.

The reaction mixture is concentrated to dryness and purified on silica to yield to 74 mg of N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-2-methylpropyl}-2-(difluoromethyl)benzamide (52%).

Mass spectrum: $[M+1] = 407$.

Example 4 : Preparation of N-(1-{[3-chloro-5-(trifluoromethyl)-2-pyridinyl]methyl}propyl)-2-(trifluoromethyl)benzamide (compound 47)

100 mg of 1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-2-butanamine (0.0004 mol), 58 μ L of 2-(trifluoromethyl)benzoyl chloride (0.0004 mol), 55 μ L of triethylamine (0.0004 mol) are stirred in 2 mL of dichloromethane at room temperature overnight.

The reaction mixture is concentrated to dryness and purified on silica to yield to 96 mg of N-(1-{[3-chloro-5-(trifluoromethyl)-2-pyridinyl]methyl}propyl)-2-(trifluoromethyl)benzamide (78 %).

Mass spectrum: $[M+1] = 425$.

Example 5 : Preparation of N-{2-(acetylamino)-2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-(trifluoromethyl)benzamide (compound 48)

108 mg of N-{2-amino-1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}acetamide trifluoroacetate (0.0003 mol), 76 mg of 2-(trifluoromethyl)benzoic acid (0.0003 mol), 83 mg of 4-(4,6-dimethoxy[1.3.5]triazin-2-yl)-4-methylmorpholinium chloride hydrate (0.0003 mol) and 41 μ L of triethylamine (0.0003 mol) are stirred in 3 mL of ethanol at room temperature overnight.

The reaction mixture is concentrated to dryness and purified on silica to yield to 23.5 mg of N-(1-{[3-chloro-5-(trifluoromethyl)-2-pyridinyl]methyl}propyl)-2-(trifluoromethyl)benzamide.

Mass spectrum: $[M+1] = 473$.

Example 6 : Preparation of N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-1-phenylethyl}-2-(trifluoromethyl)benzamide (compound 49)

79 mg of 2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-1-phenylethanamine
 5 (0.0003 mol), 57 mg of 2-(trifluoromethyl)benzoic acid (0.0003 mol), 83
 mg of 4-(4,6-dimethoxy[1.3.5]triazin-2-yl)-4-methylmorpholinium chloride hydrate
 (0.0003 mol) are stirred in 3 mL of ethanol at room temperature overnight.

The reaction mixture is concentrated to dryness and purified on silica to yield
 10 to 45 mg of N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-1-phenylethyl}-2-
 (trifluoromethyl)benzamide.

Mass spectrum: $[M+1] = 454$.

*Examples of process for the preparation of starting material (intermediates of
 general formula II) :*

15

Example 7 : Preparation of 2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-1-phenylethanamine

Preparation of 2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-1-phenylethanone

20 To a suspension of 2.6 g (0.065 mol) of sodium hydride 60% in
 dimethoxyethane at room temperature is added 3.4 mL (0.029 mol) of acetophenone.
 After 45 min., 5.55 mL (0.038 mol) of 2,3-dichloro-5-(trifluoromethyl)pyridine is
 added. After 25 min., the reaction mixture is poured over 100mL of hydrochloric
 acid 1N, extracted twice with 100mL of ethyl acetate.

25 The organic phase is washed twice 100 mL of water, dried over magnesium
 sulfate, filtered and concentrated to provide 15g of crude material which is purified
 over a column of silica by using a mixture of heptane and ethyl acetate as eluent, to
 yield to 5.74 g of desired product 2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-1-
 phenylethanone (74%).

30 RMN ^1H δ (ppm) 8,73 ; (1H, s) ; 7,95 (1H, s) ; 7,45 (2H, m) ; 7,42 (2H, m) ; 4,75
 (2H, s).

Preparation of 2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-1-phenylethanamine

35 5.6g (0.0187 mol) of 2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-1-
 phenylethanone are diluted in 50mL of methanol. 50 g of molecular sieves 3 Å ,
 14.4g (0.187 mol) of ammonium acetate and 2.45g (0.037 mol) of sodium

cyanoborohydride are then consequently added. The pH is adjusted to 5-6 with acetic acid (1 mL). After of 4 days of reaction at room temperature, the medium is filtered and 1M aqueous sodium hydroxide is added until pH = 12. 150 mL of ethyl acetate are added, after separation, the aqueous phase is extracted twice with 150 mL of ethyl acetate.

The organic phase is washed with 100 mL of brine, 100 mL of water; dried over magnesium sulfate, filtered and concentrated to yield to 1.3g of desired product 2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-1-phenylethanamine (23%).
Mass spectrum: $[M+1] = 301$.

Example 8 : Preparation of 2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-1-butanamine hydrochloride

Preparation of methyl [3-chloro-5-(trifluoromethyl)-2-pyridinyl](cyano)acetate

Under argon, 116 g of sodium hydride 60% in dispersion in oil (2.91 mol, 1.8 eq.) was suspended in 3L of DMF. The suspension was cooled down in an ice-water bath. 160 g (1.616 mol, 1.0 eq) of methyl cyanoacetate in solution in 200 mL of DMF was added dropwise under stirring. Temperature rose to 50°C and hydrogen was released. Once bubbling was over, 350g (1.616 mol, 1.0 eq) of 2,3-dichloro-5-(trifluoromethyl)-pyridine was added under stirring. The mixture was stirred overnight at room temperature. 50mL of methanol were added dropwise to quench the reaction. The reaction mixture was poured in 5L of water. pH was adjusted to 3-4 with concentrated hydrochloric acid.

The yellow precipitate which formed was filtered and washed thoroughly with water and pentane. 414g of methyl [3-chloro-5-(trifluoromethyl)-2-pyridinyl](cyano)acetate (92%) were recovered.
Mass spectrum: $[M+1] = 279$.

Preparation of [3-chloro-5-(trifluoromethyl)-2-pyridinyl] acetonitrile

314g (1.13 mol, 1eq.) of methyl [3-chloro-5-(trifluoromethyl)-2-pyridinyl](cyano)acetate and 22g (0.38mol, 0.33 eq.) of sodium chloride were dissolved in mixture of 44mL of water and 1.1L of DMSO. The reaction mixture was stirred and heated to 160°C. Gases were released, once the bubbling was over, the reaction was allowed to come back to room temperature under stirring. 1L of water and 0.5L of dichloromethane were added. After separation, the aqueous phase was extracted twice with 0.5L of DCM.

The organic phase was extracted twice with 0.5L of water and dried over magnesium sulfate. After concentration, the crude product was diluted in 100mL of DCM and eluted with EtOAc / Heptane (20/80) over a bed of silica. The filtrate was concentrated to yield 227g of [3-chloro-5-(trifluoromethyl)-2-pyridinyl] acetonitrile (91%).

Mass spectrum: $[M+1] = 223$.

Preparation of 2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]butanenitrile

199g (0.9 mol) of 3 of [3-chloro-5-(trifluoromethyl)-2-pyridinyl] acetonitrile are dissolved in 3L of THF at -5°C . A solution of 106g (0.945 mol) of potassium *tert*-butoxide in 0.6 L of THF is slowly added to the reaction medium. After 2H, 147g (0.945 mol) of ethyl iodide is added dropwise to the reaction mixture which is stirred overnight at room temperature. 3L of water and 2L of ethyl acetate are added to the reaction mixture, after separation, the aqueous phase is extracted twice with 500 mL of ethyl acetate.

The organic phase is washed with 4L of brine, 1L of water, dried over magnesium sulfate, concentrated to yield to 223g of desired product 2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]butanenitrile (100%).

RMN ^1H δ (ppm) 8,7 (1H, d, $J = 1,5\text{Hz}$) ; 7,91 (1H, d, $J = 1,5\text{Hz}$) ; 4,37 (1H, dd, $J = 6,57\text{Hz} - 7,83\text{Hz}$) ; 2,00 (2H, m) ; 1,06 (3H, t, $J = 7.33\text{Hz}$).

Preparation of *tert*-butyl 2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]butylcarbamate

189 g of 2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]butanenitrile (0.758 mol), 331g (1.51 mol) of *di**tert*-butyl carbonate, 198 g (0.834 mol) of nickel chloride (II) - hexahydrate are stirred in 2 L of methanol at room temperature. 200.6 g (5.31 mol) of sodium borohydride are added portionwise. After 3 hours of stirring, the reaction mixture is filtered over Supercel, 2 L of ethyl acetate are added, followed by 1.5 L of an aqueous solution of sodium bicarbonate.

After separation, the organic phase is washed thrice with 1 L of an aqueous solution of sodium bicarbonate, dried over magnesium sulfate, concentrated to yield to 375 g of crude material which is purified over silica to produce 147 g of desired product *tert*-butyl 2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]butylcarbamate (50,5%).

RMN ^1H δ (ppm) 8,88 (1H, d, $J = 1,5\text{Hz}$) ; 8,04 (1H, d, $J = 1,5\text{Hz}$) ; 5,04 (1H, b) ; 3,78 (1H, m) ; 3,67 (2H, m) ; 1,88 (2H, m) ; 1,34 (9H, s) ; 1,01 (3H, t, $J = 7.33\text{Hz}$).

Preparation of 2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-1-butanamine hydrochloride

146 g of tert-butyl 2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]butylcarbamate (0.414 mol) are dissolved in 2 L of dichloromethane. 190 mL of trifluoroacetic acid are added dropwise. The reaction mixture is stirred at room temperature for twelve hours, concentrated to dryness and diluted in 300 mL of hydrochloric acid 2M.

After 2 H, the reaction mixture is concentrated to dryness to provide 104 g of desired product 2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-1-butanamine hydrochloride (87%).
m.p. = 139-142°C.

Example 9 : Preparation of 1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-2-butanamine

Preparation of methyl 2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-3-oxopentanoate

4.45 g of potassium hydroxide 85% are suspended in 40 mL of DMF, 4.37 g of methyl 3-oxopentanoate (0.036 mol) are subsequently added. The reaction mixture is warmed to 50°C and 6 mL of 2,3-dichloro-5-(trifluoromethyl)pyridine are introduced. The reaction mixture is stirred at 50°C for 4 hours, quenched with 150 mL of an aqueous solution of NaH_2PO_4 (1M), extracted thrice with 150 mL of ethyl acetate.

The organic phase is washed with 150 mL of brine, 150 mL of water, dried over magnesium sulfate and concentrated to provide 8.8 g of crude material which is purified on silica to yield 2.09 g of desired product methyl 2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-3-oxopentanoate (24%).

Mass spectrum: $[M+1] = 310$.

Preparation of 1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-2-butanone

2.08 g of methyl 2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-3-oxopentanoate (0.00677 mol) and 0.12 g of sodium chloride are dissolved in a mixture of water (0.25 mL) and dimethylsulfoxide (20 mL). The reaction medium is stirred at a temperature of 130°C for 8 hours. After cooling, 100 mL of water are added to the reaction mixture which is extracted twice with 150 mL of ethyl acetate.

The organic phase is washed with twice 100 mL of water, dried over magnesium sulfate and concentrated. After a purification on silica, 0.67 g of desired product 1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-2-butanone are produced (39%). Mass spectrum: $[M+1] = 253$.

5

Preparation of 1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-2-butanamine

0.64 g of 1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-2-butanone (0.0025 mol) are diluted in 5 mL of methanol. 7.0 g of molecular sieves 3 Å, 1.90 g (0.025 mol) of ammonium acetate and 0.39 g (0.0052 mol) of sodium cyanoborohydride are then consequently added. The reaction medium is stirred overnight at room temperature. After filtration, the pH is adjusted to 9 with an aqueous solution of sodium hydroxide 1 M. The reaction mixture is concentrated to dryness. 30 mL of ethyl acetate are added. The organic phase is washed with an aqueous solution of sodium hydroxide 1 M, brine, and water; dried over magnesium sulfate, filtered and concentrated to dryness. The crude material is dissolved in 15 mL of hydrochloric acid 1M, extracted with 15 mL of ethyl acetate.

10

The aqueous phase is then basified with an aqueous solution of sodium hydroxide 1 M, extracted thrice with 15 mL of ethyl acetate, dried over magnesium sulfate, filtered and concentrated to dryness to provide 0.21 g of desired product 1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-2-butanamine (32%).
Mass spectrum : $[M+1] = 253$.

15

20

Example 10 : Preparation of N-{2-amino-1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}acetamide trifluoroacetate

25

Preparation of [3-chloro-5-(trifluoromethyl)-2-pyridinyl]((diphenylmethylene)amino)acetonitrile

To a suspension of sodium hydride 60% (4.0g, 0.1 mol) in dimethylformamide (50 mL) at -10°C was dropwise added a solution of N-(diphenylmethylene)aminoacetonitrile (11.1 g, 0.05 mol) 60 mL of dimethylformamide. After one hour of stirring is added to the reaction mixture, 7 mL of 2,3-dichloro-5-(trifluoromethyl)pyridine (0.052 mol) are added to the reaction mixture. The reaction medium is allowed to room temperature, stirred for 3 hours, quenched with a mixture of diethyl ether (500 mL) and an aqueous solution of ammonium chloride 10% (500mL). After separation, the organic phase is dried over magnesium sulfate, filtered and concentrated.

30

35

The crude material is purified on silica to yield to 17.8g of [3-chloro-5-(trifluoromethyl)-2-pyridinyl][(diphenylmethylene)amino]acetonitrile (90%).
m.p. = 105-108°C.

5 Preparation of amino[3-chloro-5-(trifluoromethyl)-2-pyridinyl]acetonitrile hydrochloride

15.0 g of [3-chloro-5-(trifluoromethyl)-2-pyridinyl][(diphenylmethylene)amino]acetonitrile (0.037 mol) are stirred at room temperature for 3h in a mixture of dichloromethane (15 mL) and hydrochloric acid 10% (15 mL). Phases are separated and the organic phase is washed with 15 mL of hydrochloric acid 10%.

The aqueous phase is concentrated to dryness to provide a pink solid which is washed with diethyl ether, filtered and dried to provide 8.10 g of desired product amino[3-chloro-5-(trifluoromethyl)-2-pyridinyl]acetonitrile hydrochloride (79 %).
15 m.p. = 258-260°C.

Preparation of N-[[3-chloro-5-(trifluoromethyl)-2-pyridinyl](cyano)methyl]acetamide

0.67 g of amino[3-chloro-5-(trifluoromethyl)-2-pyridinyl]acetonitrile hydrochloride (0.00245 mol) are dissolved in 15 mL of dichloromethane, 0.68 mL of triethylamine are added . After 10 min. at room temperature, 0.18 mL of acetyl chloride (0.00245 mol) are added, the reaction mixture is stirred overnight. The reaction is quenched with 20 mL of water.

After separation, the organic phase is washed with 20 mL of water, dried on magnesium sulfate, filtered and concentrated to dryness to produce 0.75 g of crude material which is purified on silica to yield 0.54 g of desired product N-[[3-chloro-5-(trifluoromethyl)-2-pyridinyl](cyano)methyl]acetamide (80%).
25 Mass spectrum: [M+1] = 278.

30 Preparation of tert-butyl 2-(acetylamino)-2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethylcarbamate

5.00 g of N-[[3-chloro-5-(trifluoromethyl)-2-pyridinyl](cyano)methyl]acetamide (0.018 mol), 7.86 g (0.036 mol) of di-tert-butyl carbonate, 4.28 g (0.018 mol) of nickel chloride (II) - hexahydrate are stirred in 25mL of methanol at room temperature. 3.40 g (0.09 mol) of sodium borohydride are added portionwise. The reaction mixture is stirred overnight. 200 mL of ethyl acetate

are added, followed by 50 mL of water. After separation, the aqueous phase is extracted twice 50 mL of ethyl acetate.

The organic phase is dried over magnesium sulfate, concentrated to yield to 4.01 g of crude material which is purified over silica to produce 1.35 g of desired product tert-butyl 2-(acetylamino)-2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethylcarbamate (19%).

Mass spectrum: $[M+1] = 382$.

Preparation of N-{2-amino-1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}acetamide trifluoroacetate

1.30 g of tert-butyl 2-(acetylamino)-2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethylcarbamate (0.034 mol) are diluted in a mixture of dichloromethane (2.5 mL) and trifluoroacetic acid (2.5 mL). After stirring overnight at room temperature, the reaction mixture is concentrated to dryness to yield to 1.34 g of desired product N-{2-amino-1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}acetamide trifluoroacetate (99 %).

Mass spectrum: $[M+1-HCl] = 282$.

Example 11 : Preparation of 2-amino-1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethanol

Preparation of 1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl] ethanone

To 225 ml of dry toluene were added 210 ml (0.29 mol) of a 1.4 M solution of methylmagnesium bromide in toluene/tetrahydrofuran 75:25. The solution was cooled to -5°C and 30 g (0.145 mol) of 3-chloro-5-(trifluoromethyl)-2-pyridinecarbonitrile, were slowly added in 2 hours at 0°C . After addition, the dark solution was further stirred at room temperature for 5 hours. The reaction mixture was neutralized by 350 ml of 1N hydrochloric acid and stirred 3 hours at room temperature. The aqueous phase was then reextracted by ethyl acetate (3 x 200 ml), washed with water (300 ml) and dried over magnesium. The solvent was evaporated under reduced pressure to give 33.2 g of the crude product as a brown oil.

The crude product was purified by flash chromatography on silica gel (eluent: heptane/ethyl acetate 9:1) to give 1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethanone: 13.1 g (40 %) as a yellow oil.

Mass spectrum: $[M+1] = 224$.

Preparation of 2-bromo-1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl] ethanone

To a solution of 16.6 g (0.074 mol) of 1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl] ethanone in 150 ml of dry tetrahydrofuran, were added portionwise at room temperature 27.8 g (0.074 mol) of phenyltrimethylammonium tribromide. The solution was stirred for 3 hours at room temperature. The formed solid was removed by filtration and the mother liquors were concentrated under vacuum. The resulting orange oil (32.7 g) was purified by flash chromatography on silica gel (eluent: heptane/dichloromethane 3:1) to give 2-bromo-1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethanone: 10.4 g (46 %) as a yellow oil.

Mass spectrum: $[M-1] = 302$.

Preparation of 2-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-2-oxoethyl}-1H-isoindole-1,3(2H)-dione

To a solution of 1 g (3.3 mmol) of 2-bromo-1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl] ethanone in 10 ml of 2-butanone were successively added 50 mg (0.33 mmol) of potassium iodide and 1.23 g (6.6 mmole) of potassium phthalimide. The reaction mixture was stirred at 86 °C for 2 hours. The reaction mixture was poured into 100 ml of water and the aqueous phase was extracted with ethyl acetate (2 x 50 ml) and the organic phase was washed with water and dried over magnesium sulfate.

The solvent was evaporated under reduced pressure and the resulting solid was triturated in dichloromethane to give 2-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-2-oxoethyl}-1H-isoindole-1,3(2H)-dione: 0.35 g (29 %) as a beige solid ; mp = 162 °C.

Mass spectrum: $[M+1] = 369$.

Preparation of 2-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-2-hydroxyethyl}-1H-isoindole-1,3(2H)-dione

To a suspension of 36 mg (0.95 mmol) of sodium borohydride in 5 ml of methanol, was added at 0°C, 700 mg (1.9 mmol) of 2-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-2-oxoethyl}-1H-isoindole-1,3(2H)-dione. The reaction mixture was stirred at room temperature for 1 hour. The pH was brought to 7 by addition of 1N hydrochloric acid and methanol was removed under reduced pressure. The residue was reextracted by dichloromethane (2 x 50 ml) and the organic phase was washed with brine and dried over magnesium sulfate. The solvent was evaporated to give 0.55 g of the crude product as a yellow solid.

The crude product was purified by flash chromatography on silica gel (eluent: heptane/ethyl acetate 8:2) to give 2-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-2-hydroxyethyl}-1*H*-isoindole-1,3(2*H*)-dione: 0.3 g (42 %) as a cream-coloured solid. Mass spectrum: $[M+1] = 371$.

5

Preparation of 2-amino-1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethanol

To a suspension of 150 mg (0.4 mmol) of 2-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-2-hydroxyethyl}-1*H*-isoindole-1,3(2*H*)-dione in 3 ml of ethanol, was added 0.02 ml (0.4 mmol) of hydrazine monohydrate.

10

The reaction mixture was heated at reflux for 4 hours. After cooling, the white solid was filtered off and the mother liquors were concentrated under vacuum to give 2-amino-1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethanol: 50 mg (52 %) as a yellow solid.

Mass spectrum: $[M+1] = 241$.

15

Examples of biological activity of the compound of general formula (I)

Example A : *in vivo* test on *Alternaria brassicae* (Leaf spot of crucifers)

5

The active ingredient tested is prepared by potter homogenisation in a concentrated suspension type formulation at 100 g/l. This suspension is then diluted with water to obtain the desired active material concentration.

10 Radish plants (Pernot variety) in starter cups, sown on a 50/50 peat soil-pozzolana substrate and grown at 18-20°C, are treated at the cotyledon stage by spraying with the aqueous suspension described above.

Plants, used as controls, are treated with an aqueous solution not containing the active material.

15 After 24 hours, the plants are contaminated by spraying them with an aqueous suspension of *Alternaria brassicae* spores (40,000 spores per cm³). The spores are collected from a 12-13-day-old culture.

The contaminated radish plants are incubated for 6-7 days at about 18°C, under a humid atmosphere.

20 Grading is carried out 6 to 7 days after the contamination, in comparison with the control plants.

Under these conditions, good (at least 50%) to total protection is observed at a dose of 330ppm with the following compounds : 3, 4, 7, 8, 9, 10, 11, 12, 13, 15, 19, 21, 23, 25, 32, 33, 35, 36, 37, 38, 42, 43, 47.

25 Example B : *in vivo* test on *Erysiphe graminis* f. sp. *tritici* (powdery mildew of wheat)

The active ingredient tested is prepared by potter homogenisation in a concentrated suspension type formulation at 100 g/l. This suspension is then diluted
30 with water to obtain the desired active material concentration.

Wheat plants (Audace variety) in starter cups, sown on 50/50 peat soil-pozzolana substrate and grown at 12°C, are treated at the 1-leaf stage (10 cm tall) by spraying with the aqueous suspension described above.

35 Plants, used as controls, are treated with an aqueous solution not containing the active material.

After 24 hours, the plants are contaminated by dusting them with *Erysiphe graminis* f. sp. *tritici* spores, the dusting being carried out using diseased plants.

Grading is carried out 7 to 14 days after the contamination, in comparison with the control plants.

5 Under these conditions, good (at least 50%) total protection is observed at a dose of 330 ppm with the following compounds : 32, 33, 38, 40, 43.

Example C : *in vivo* test on *Pyrenophora teres* (Barley Net blotch).

10 The active ingredient tested is prepared by potter homogenisation in a concentrated suspension type formulation at 100 g/l. This suspension is then diluted with water to obtain the desired active material concentration.

Barley plants (Express variety) in starter cups, sown on a 50/50 peat soil-pozzolana substrate and grown at 12°C, are treated at the 1-leaf stage (10 cm
15 tall) by spraying with the aqueous suspension described above. Plants, used as controls, are treated with an aqueous solution not containing the active material.


After 24 hours, the plants are contaminated by spraying them with an aqueous suspension of *Pyrenophora teres* spores (12,000 spores per ml). The spores are collected from a 12-day-old culture. The contaminated barley plants are incubated
20 for 24 hours at about 20°C and at 100% relative humidity, and then for 12 days at 80% relative humidity.

Grading is carried out 12 days after the contamination, in comparison with the control plants.

25 Under these conditions, good (at least 50%) total protection is observed at a dose of 330 ppm with the following compounds : 7, 8, 9, 10, 13, 30, 32, 33, 37, 38, 39, 40, 41, 43, 44, 47.

CLAIMS

- 5 1. A compound of general formula (I) :



in which :

- n is 1, 2, or 3;
- p is 1, 2, 3 or 4;
- R^a is a C₁-C₆-halogenoalkyl having 1 to 5 halogen atoms;
- each substituent X is chosen, independently of the others, as being a hydrogen atom, a halogen atom, a C₁-C₆-alkyl or a C₁-C₆-halogenoalkyl;
- R¹, R², R³ and R⁴ are chosen, independently of the others as being a hydrogen atom, a halogen atom, a cyano group, a hydroxy group, an amino group, a sulfanyl group, a formyl group, a formyloxy group, a formylamino group, a carboxy group, a carbamoyl group, a N-hydroxycarbamoyl group, a carbamate group, a (hydroxyimino)-C₁-C₆-alkyl group, a C₁-C₆-alkyl, a C₂-C₆-alkenyl, a C₂-C₆-alkynyl, a C₁-C₆-alkylamino, a di-C₁-C₆-alkylamino, a C₁-C₆-alkoxy, a C₁-C₆-halogenoalkyl having 1 to 5 halogen atoms, a C₁-C₆-halogenoalkoxy having 1 to 5 halogen atoms, a C₁-C₆-alkylsulfanyl, a C₁-C₆-halogenoalkylsulfanyl having 1 to 5 halogen atoms, a C₂-C₆-alkenyloxy, a C₂-C₆-halogenoalkenyloxy having 1 to 5 halogen atoms, a C₃-C₆-alkynyloxy, a C₃-C₆-halogenoalkynyloxy having 1 to 5 halogen atoms, a C₃-C₆-cycloalkyl, a C₃-C₆-halogenocycloalkyl having 1 to 5 halogen atoms, a C₁-C₆-alkylcarbonyl, a C₁-C₆-halogenoalkylcarbonyl having 1 to 5 halogen atoms, a C₁-C₆-alkylcarbamoyl, a di-C₁-C₆-alkylcarbamoyl, a N-C₁-C₆-alkyloxycarbamoyl, a C₁-C₆-alkoxycarbamoyl, a N-C₁-C₆-alkyl-C₁-C₆-alkoxycarbamoyl, a C₁-C₆-alkoxycarbonyl, a C₁-C₆-halogenoalkoxycarbonyl having 1 to 5 halogen atoms, a C₁-C₆-alkylcarbonyloxy, a C₁-C₆-halogenoalkylcarbonyloxy having 1 to 5 halogen atoms, a C₁-C₆-alkylcarbonylamino, a C₁-C₆-halogenoalkylcarbonylamino having 1 to 5 halogen atoms, a C₁-C₆-alkylaminocarbonyloxy, a di-C₁-C₆-alkylaminocarbonyloxy, a C₁-C₆-alkyloxycarbonyloxy, a C₁-C₆-alkylsulphenyl, a C₁-C₆-

halogenoalkylsulphenyl having 1 to 5 halogen atoms, a C₁-C₆-alkylsulphinyl, a C₁-C₆-halogenoalkylsulphinyl having 1 to 5 halogen atoms, a C₁-C₆-alkylsulphonyl, a C₁-C₆-halogenoalkylsulphonyl having 1 to 5 halogen atoms, a benzyl, a benzyloxy, a benzylsulfanyl, a benzylsulfinyl, a benzylsulfonyl, a benzylamino, a phenoxy, a phenylsulfanyl, a phenylsulfinyl, a phenylsulfonyl, a phenylamino, a phenylcarbonylamino, a 2,6 dichlorophenyl-carbonylamino group or a phenyl group,

with the proviso that when three of the four substituents R¹, R², R³ and R⁴ are a hydrogen atom, then the fourth substituent is not a hydrogen atom;

- R⁵ is chosen as being a hydrogen atom, a cyano group, a formyl group, a hydroxy group, a C₁-C₆-alkyl, a C₁-C₆-halogenoalkyl having 1 to 5 halogen atoms, a C₁-C₆-alkoxy, a C₁-C₆-halogenoalkoxy having 1 to 5 halogen atoms, a C₃-C₆-halogenocycloalkyl having 1 to 5 halogen atoms, a C₂-C₆-alkenyl, a C₂-C₆-alkynyl, a C₁-C₆-alkoxy-C₁-C₆-alkyl, a C₁-C₆-cyanoalkyl, a C₁-C₆-aminoalkyl, a C₁-C₆-alkylamino-C₁-C₆-alkyl, a di-C₁-C₆-alkylamino-C₁-C₆-alkyl, a C₁-C₆-alkylcarbonyl, a C₁-C₆-halogenalkylcarbonyl having 1 to 5 halogen atoms, a C₁-C₆-alkyloxycarbonyl, a C₃-C₇-cycloalkyl, a C₃-C₇-halogenocycloalkyl having 1 to 5 halogen atoms, a C₃-C₇-cycloalkyl-C₁-C₆-alkyl, a C₁-C₆-benzyloxycarbonyl, a C₁-C₆-alkoxy-C₁-C₆-alkylcarbonyl, a C₁-C₆-alkylsulfonyl or a C₁-C₆-halogenoalkylsulfonyl having 1 to 5 halogen atoms;

- Y is the same or different and is a hydrogen atom, a halogen atom, a nitro group, a cyano group, a hydroxy group, an amino group, a sulfanyl group, a pentafluoro-λ⁶-sulfanyl group, a formyl group, a formyloxy group, a formylamino group, a carboxy group, a C₁-C₈-alkyl, a C₁-C₈-halogenoalkyl having 1 to 5 halogen atoms, a C₂-C₈-alkenyl, a C₂-C₈-alkynyl, a C₁-C₈-alkylamino, a di-C₁-C₈-alkylamino, a C₁-C₈-alkoxy, a C₁-C₈-halogenoalkoxy having 1 to 5 halogen atoms, a C₁-C₈-alkoxy-C₂-C₈-alkenyl, a C₁-C₈-alkylsulfanyl, a C₁-C₈-halogenoalkylsulfanyl having 1 to 5 halogen atoms, a C₁-C₈-alkoxycarbonyl, a C₁-C₈-halogenoalkoxycarbonyl having 1 to 5 halogen atoms, a C₁-C₈-alkylcarbonyloxy, a C₁-C₈-halogenoalkylcarbonyloxy having 1 to 5 halogen atoms, a C₁-C₈-alkylsulphenyl, a C₁-C₈-halogenoalkylsulphenyl having 1 to 5 halogen atoms, a C₁-C₈-alkylsulphinyl, a C₁-C₈-halogenoalkylsulphinyl having 1 to 5 halogen atoms, a C₁-C₈-alkylsulphonyl, a C₁-C₈-halogenoalkylsulphonyl having 1 to 5 halogen atoms or a C₁-C₈-alkylsulfonamide; and


- R^b is a halogen atom, a nitro group, a cyano group, an amino group, a sulfanyl group, a pentafluoro-λ⁶-sulfanyl group, a formyl group, a formyloxy group, a formylamino group, a carboxy group, a C₁-C₆-alkyl, a C₁-C₆-halogenoalkyl having

1 to 5 halogen atoms, a C₂-C₆-alkenyl, a C₂-C₆-alkynyl, a C₁-C₆-alkylamino, a di-C₁-C₆-alkylamino, a C₁-C₆-alkoxy, a C₁-C₆-halogenoalkoxy having 1 to 5 halogen atoms, a C₁-C₆-alkoxy-C₂-C₆-alkenyl, a C₁-C₆-alkylsulfanyl, a C₁-C₆-halogenoalkylsulfanyl having 1 to 5 halogen atoms, a C₁-C₆-alkoxycarbonyl, a C₁-C₆-halogenoalkoxycarbonyl having 1 to 5 halogen atoms, a C₁-C₆-alkylcarbonyloxy, a C₁-C₆-halogenoalkylcarbonyloxy having 1 to 5 halogen atoms, a C₁-C₆-alkylsulphenyl, a C₁-C₆-halogenoalkylsulphenyl having 1 to 5 halogen atoms, a C₁-C₆-alkylsulphanyl, a C₁-C₆-halogenoalkylsulphanyl having 1 to 5 halogen atoms, a C₁-C₆-alkylsulphonyl, a C₁-C₆-halogenoalkylsulphonyl having 1 to 5 halogen atoms or a C₁-C₆-alkylsulfonamide;


as well as its salts, N-oxydes, metallic complexes, metalloidic complexes and optically active isomers.

2. A compound according to claim 1, characterised in that n is 1 or 2.
3. A compound according to claim 1 or 2, characterised in that X is a halogen atom.
4. A compound according to claim 3, characterised in that X is chlorine.
5. A compound according to any of the claims 1 to 4, characterised in that R^a is -CF₃.
6. A compound according to any of the claims 1 to 5, characterised in that the 2-pyridyl is substituted in 3- and/or in 5-position.
7. A compound according to claim 6, characterised in that the 2-pyridyl is substituted in 3-position by X and in 5-position by R^a.
8. A compound according to any of the claims 1 to 7, characterised in that the 2-pyridyl is substituted in 3-position by -Cl and in 5-position by -CF₃.
9. A compound according to any of the claims 1 to 8, characterised in that R^b is a halogen atom, a C₁-C₆-alkyl, a C₁-C₆-alkoxy or a C₁-C₆-halogenoalkyl having 1 to 5 halogen atoms.

10. A compound according to any of the claims 1 to 9, characterised in that p is 1.
11. A compound according to any of the claims 1 to 10, characterised in that Y is
5 a hydrogen atom, a halogen atom or a C₁-C₆-alkyl.
12. A compound according to any of the claims 1 to 11, characterised in that R¹
and R² are chosen, independently of each other, as being a hydrogen atom, a halogen
atom, a cyano group, a hydroxy group, a C₁-C₆-alkyl, a C₁-C₆-halogenoalkyl having
10 1 to 5 halogen atoms, a C₂-C₆-alkenyl, a C₁-C₆-alkoxy, a C₁-C₆-alkylsulfanyl, a C₁-
C₆-alkylsulfenyl, a C₁-C₆-alkylsulfinyl, a C₁-C₆-alkoxycarbonyl, a C₁-C₆-
alkylcarbonylamino, a C₁-C₆-alkoxycarbonyloxy, a C₁-C₆-alkoxycarbonylamino or a
phenyl group.
13. A compound according to claim 12, characterised in that R¹ and R² are
15 chosen, independently of each other, as being a halogen atom, a C₁-C₆-alkyl, a C₁-
C₆-halogenoalkyl having 1 to 5 halogen atoms, a C₁-C₆-alkylcarbonylamino.
14. A compound according to any of the claims 1 to 13, characterised in that R³
20 and R⁴ are chosen, independently of each other, as being a hydrogen atom, a halogen
atom, a cyano group, a C₁-C₆-alkyl, a C₁-C₆-halogenoalkyl having 1 to 5 halogen
atoms, a C₁-C₆-alkylcarbonylamino or a phenyl group.
15. A compound according to claim 14, characterised in that R³ and R⁴ are
25 chosen, independently of each other, as being a halogen atom, a C₁-C₆-alkyl, a C₁-
C₆-halogenoalkyl having 1 to 5 halogen atoms or a phenyl group.
16. A compound according to any of the claims 1 to 13, characterised in that R⁵
is a hydrogen atom or a C₃-C₇-cycloalkyl.
30
17. A process for the preparation of a compound of general formula (I) as defined
in any of the claims 1 to 16, which comprises reacting a 2-pyridine derivative of
general formula (II) or one of its salt :




in which X, n, R^a, R¹, R², R³, R⁴ and R⁵ are as in any of the preceding claims;
with a carboxylic acid derivative of the general formula (III)




in which :

- Y, p and R^b are as defined in any of the preceding claims ; and
- L² is a leaving group chosen as being a halogen atom, a hydroxyl group, -OR⁶, -OCOR⁶, R⁶ being a C₁-C₆ alkyl, a C₁-C₆ haloalkyl, a benzyl, 4-methoxybenzyl, pentafluorophenyl or a group of formula



in the presence of a catalyst and, if L² is a hydroxyl group, in the presence of a condensing agent.

18. A process according to claim 17, characterised in that R⁵ is a hydrogen atom and that the process is completed by a further step according to the following reaction scheme :



in which : - R¹, R², R³, R⁴, R^a, R^b, X, Y, n and p are as defined in any of the claims 1 to 15;

- L⁵ is a leaving group chosen as being a halogen atom, a 4-methyl phenylsulfonyloxy or a methylsulfonyloxy;


comprising the reaction of a compound of general formula (Id) with a compound of general formula (XXII) to provide a compound of general formula (I).

19. A fungicidal composition comprising an effective amount of a compound according to any of the claims 1 to 16 and an agriculturally acceptable support.

20. A method for preventively or curatively combating the phytopathogenic fungi
5 of crops, characterised in that an effective and non-phytotoxic amount of a composition according to claim 19 is applied to the plant seeds or to the plant leaves and/or to the fruits of the plants or to the soil in which the plants are growing or in which it is desired to grow them.

ABSTRACT

A compound of general formula (I) :



5

A process for preparing this compound.

A fungicidal composition comprising a compound of general formula (I).

A method for treating plants by applying a compound of general formula (I) or a composition comprising it.

PCT/EP2004/014899^{SP2}



**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☒ FADED TEXT OR DRAWING

☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☒ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☒ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.